

Learning from Neighbours about a Changing State

KRISHNA DASARATHA

Boston University

BENJAMIN GOLUB

Northwestern University

and

NIR HAK

Uber Technologies, Inc.

First version received January 2020; Accepted November 2022 (Eds.)

Agents learn about a changing state using private signals and their neighbours' past estimates of the state. We present a model in which Bayesian agents in equilibrium use neighbours' estimates simply by taking weighted sums with time-invariant weights. The dynamics thus parallel those of the tractable DeGroot model of learning in networks, but arise as an equilibrium outcome rather than a behavioural assumption. We examine whether information aggregation is nearly optimal as neighbourhoods grow large. A key condition for this is *signal diversity*: each individual's neighbours have private signals that not only contain independent information, but also have sufficiently different distributions. Without signal diversity—*e.g.* if private signals are i.i.d.—learning is suboptimal in all networks and highly inefficient in some. Turning to social influence, we find it is much more sensitive to one's signal quality than to one's number of neighbours, in contrast to standard models with exogenous updating rules.

Key words: Social learning, Networks, DeGroot learning, Information aggregation, Network centrality

1. INTRODUCTION

People learn from others about conditions relevant to decisions they have to make. In many cases, the conditions—*e.g.* the state of a labour market facing workers, or the business environment relevant to an organisation—are changing. Thus, welfare depends not on learning a static “state of the world,” but rather on staying up to date with a changing state. The phenomenon of adaptation and responsiveness to new information is central in many economic applications, including in economic development, organisational behavior, and financial decision-making. When is a group of agents successful, collectively, in adapting efficiently to a changing environment? The answers lie partly in the structure of the social networks that shape agents' social learning opportunities. Our model is designed to analyse how a group's adaptability is shaped by the properties of such networks, the inflows of information into society, and the interplay of the two.

We consider overlapping generations of agents who are interested in tracking an unobserved state that evolves over time.¹ The state is a Gaussian AR(1) process: somewhat persistent, but with constant innovations to learn about. Each agent, before making a decision, engages in social learning: she observes the actions of some members of prior generations, which reveal their estimates of the state. The social learning opportunities are embedded in a network, in that one's network position determines the *neighbourhood* of peers whom one observes. Neighbourhoods reflect geographic, cultural, organisational, or other kinds of proximity.² In addition to social information, agents also receive private signals about the current state, with Gaussian distributions that may also vary with network position; in particular, some agents may receive more precise information about the state than others.

We give some examples. When a university student begins searching for jobs, she becomes interested in the state of the relevant labour market (*e.g.* typical wages for someone like her), which naturally vary over time. She uses her own private research (a private signal) but also learns from the choices of others (*e.g.* recent graduates) who have faced a similar problem. Whom she can learn from will depend on her academic specialisation, dormitory, extracurricular activities, and so forth: she will predominantly observe predecessors who are “nearby” in these ways. Similarly, when a new cohort of professionals enters a firm such as a management consultancy or law practice, they learn about the business environment from their seniors. Who works with whom, and therefore who learns from whom, is shaped by the structure of the organisation. Beyond heterogeneity in network position, agents differ in the precision of the private signals they can access from outside the network: for example, people with quantitative training may be better placed to learn from external statistical reports.

We now detail our three main contributions. The *first* is to develop a tractable model of learning about a changing state in which Bayesian updating has a simple form: each agent forms an estimate by taking a weighted average of neighbours' estimates and her private signal. Because the environment is stationary, the weights are stationary as well. Equilibrium behaviour thus yields a dynamic paralleling the DeGroot (1974) network learning model, which is famous for its tractability, but which has been criticised for its lack of canonical foundations (Golub and Sadler, 2016; Molavi *et al.*, 2018). The weights in agents' learning rules are endogenously determined because, when each agent extracts information from neighbours' estimates, the information content of those estimates depends on the neighbours' learning rules. We characterise these weights and the distributions of behaviour in a stationary equilibrium.³ These characterisations permit the analysis of comparative statics of behaviour and welfare. The model is well-behaved computationally: equilibria can be calculated quickly in networks of thousands of nodes, which makes the framework useful for structural exercises. Finally, the basic framework and stationary equilibrium definition extend readily to accommodate various kinds of behavioural updating rules, *e.g.* ones that neglect correlations in neighbours' actions.

Our *second* contribution is to analyse when equilibrium learning facilitates good information aggregation. Here we have positive and negative results. The main positive finding is that Bayesian agents in equilibrium can achieve good aggregation under a *signal diversity* condition.

1. Cf. Banerjee and Fudenberg (2004) and Wolitzky (2018), with overlapping generations and a fixed state.

2. Sethi and Yildiz (2019) argue that, even without explicit communication costs or constraints, familiarity and shared context determine the network in which people can effectively communicate.

3. In Bayesian models of learning about a fixed state, there is a time-dependence whereby rational updating rules depend on the time elapsed since the learning process started. Studies of such models often focus on an eventual rate of learning about a fixed state. See, for instance, Molavi *et al.* (2018) and Harel *et al.* (2021). This time-dependence is absent in our stationary environment; equilibrium outcomes can be summarised by steady-state updating weights and error rates.

To formalise “good aggregation,” we first note that in our model social information is valuable to agents insofar as it allows them to estimate the state before the current period; our measure of aggregation quality is the accuracy of these estimates. We say good aggregation occurs if each agent has an estimate nearly as precise as if she simply knew the previous state. We say *signal diversity* holds if each individual has at least two different private signal precisions represented in large numbers in her neighbourhood. The positive result says that signal diversity is sufficient for good aggregation, robustly across a large class of random networks (ones arising from stochastic block models satisfying certain technical conditions). Signal diversity is valuable because it leads to diversity of neighbours’ strategies: it makes them use recent and older information differently from one another. An agent observing such neighbours, in turn, can use the diversity for statistical identification, constructing a precise estimate of the most recent state. We illustrate this key idea in an example at the end of the Introduction.

We complement the positive finding with two negative results showing that both the “signal diversity” and the “Bayesian” conditions are important. First, signal homogeneity turns out to be a fundamental obstruction to good aggregation. In an environment where everyone’s private signals are conditionally independent and identically distributed, equilibrium aggregation is bound to be inefficient. We begin by establishing this point in highly symmetric networks, where the failure of aggregation is shown to have severe welfare consequences, making each agent worse off by an unbounded amount relative to a world with signal diversity. A more general finding is that in large networks with signal homogeneity, it is impossible in equilibrium to achieve accuracies of aggregation of the same order as in our positive result under signal diversity. One might have thought that diversity of neighbours’ network positions (and thus learning opportunities) can substitute for diversity of their private signal distributions; our negative result shows that network diversity is a poor substitute for signal diversity.

We next show that the “Bayesian” condition is also important for the good-aggregation result. To do this, we contrast the learning of Bayesians best-responding to others’ learning rules with that of a naive population too unsophisticated to account for correlations in neighbours’ learning errors, as in some canonical behavioural learning models (Eyster and Rabin, 2010).⁴ We identify a class of such models in which information aggregation is essentially guaranteed to fall short of good aggregation benchmarks for all agents. The deficiencies of naive learning rules are different from and more severe than those in similar problems with an unchanging state, where naive heuristics can aggregate information very well.⁵

Our *third* contribution is to study social influence, an outcome of central importance in network theory. We define a notion of steady-state social influence—how an idiosyncratic change in an individual’s information affects others’ average behaviour. This is analogous to the definition of influence in the standard DeGroot model (where updating weights are exogenous). The endogenous determination of weights makes a big difference for how the environment affects social influence. Relative to the DeGroot model benchmark studied in DeMarzo *et al.* (2003), an agent’s social influence is much more sensitive to the quality of her private information. On the other hand, just as in the standard benchmark, an agent’s influence is approximately proportional to her degree.

4. See also Bala and Goyal (1998), a seminal model of boundedly rational learning rules in networks.

5. In analogous fixed-state environments where individuals have sufficiently many observations, if everyone uses certain simple and stationary DeGroot-style heuristics (requiring no sophistication about correlations between neighbours’ behaviour), they can learn the state precisely (Golub and Jackson, 2010; Jadbabaie *et al.*, 2012). A changing state makes such imitative heuristics quite inefficient.




FIGURE 1

The network used in the “value of diversity” example

A closing discussion makes two main points. First, some of our theoretical aggregation results use large random graphs. We perform a numerical exercise to show that the main message about information aggregation—diversity of signal types helps learning—remains valid when we calculate equilibria on graphs reflecting real social networks. Second, our analysis generalises readily to richer models of multidimensional states and signals. As one application of such a generalisation, we consider a manager who wishes to facilitate better learning in an organisation, and ask what distributions of expertise such a designer would prefer. Our results provide a distinctive rationale for *informational specialisation* as a design feature that facilitates good information aggregation.

1.1. An example

We now present a simple example that illustrates our dynamic model, highlights obstacles to learning that distinctively arise in a dynamic environment, and gives a sense of some of the main forces that play a role in our results on the quality of learning.

Consider a particular environment, with a single perfectly informed source S ; many media outlets M_1, \dots, M_n with access to the source as well as some independent private information; and the general public. The public consists of many individuals who learn only from the media outlets. We are interested in how well each member of the public could learn by following many media outlets. More precisely, we consider the example shown in Figure 1 and think of P as a generic member of the large public.

The state θ_t follows a Gaussian random walk: $\theta_t = \theta_{t-1} + v_t$, where the innovations v_t are standard normal. Each period, the source learns the state θ_t and takes an action (which can be thought of simply as making an announcement) that reveals it. The media outlets observe the source’s action from the previous period, which is θ_{t-1} . At each time period, they also receive noisy private signals, $s_{M_i,t} = \theta_t + \eta_{M_i,t}$ with normally distributed, independent, mean-zero errors $\eta_{M_i,t}$. They then announce their posterior means of θ_t , which we denote by $a_{M_i,t}$. The member of the public, in a given period t , makes an estimate based on the observations $a_{M_1,t-1}, \dots, a_{M_n,t-1}$ of media outlets’ actions in the previous period. All agents are short lived: they see actions in their neighbourhoods one period ago, and then they take an action that reveals their posterior belief of the state.

If we had an *unchanging* state but the same signals and observation structure, learning would trivially be perfect: media outlets would learn the state from the source and report it to the public.

In the dynamic environment, given that P has no signal, she can at best hope to learn θ_{t-1} (and use that to estimate θ_t). Can this benchmark be achieved, and if so, when?

A typical estimate of a media outlet at time t is a linear combination of $s_{M_i,t}$ and θ_{t-1} (the latter being the information that the media outlets learned from the source S). In particular, the estimate of media outlet M_i at time t can be expressed as

$$a_{M_i,t} = w_i s_{M_i,t} + (1 - w_i) \theta_{t-1},$$

where the weight w_i on the media outlet's signal is increasing in the precision of that signal. We give the public no private signal, for simplicity only.

Suppose first that the media outlets have identically distributed private signals. Because the member of the public observes many symmetric media outlets, it turns out that her best estimate of the state, $a_{P,t}$, is simply the average of the estimates of the media outlets. Since each of these outlets uses the same weight $w_i = w$ on its private signal, we may write

$$a_{P,t} = w \sum_{i=1}^n \frac{s_{M_i,t-1}}{n} + (1 - w) \theta_{t-2} \approx w \theta_{t-1} + (1 - w) \theta_{t-2}.$$

That is, P 's estimate is an average of media private signals from last period ($t - 1$), combined with what the media learned from the source, which tracks the state in the period before that ($t - 2$). In the approximate equality, we have used the fact that an average of many private signals is approximately equal to the state, by our assumption of independent errors. No matter how many media outlets there are, and even though each has independent information about θ_{t-1} , the public's beliefs are confounded by older information.

What if, instead, half of the media outlets (say $M_1, \dots, M_{n/2}$) have more precise private signals than the other half, perhaps because these outlets have invested more in expertise on this topic? The media outlets with more precise signals, called group A , will then place weight w_A on their private signals, while the media outlets with less precise signals (group B) use a smaller weight, w_B . We will now argue that a member of the public can extract more information from the media in this setting. In particular, she can first compute the averages of the two groups' actions

$$\begin{aligned} \text{type } A \text{ average action at time } t - 1 &= w_A \sum_{i=1}^{n/2} \frac{s_{M_i,t-1}}{n/2} + (1 - w_A) \theta_{t-2} \\ &\approx w_A \theta_{t-1} + (1 - w_A) \theta_{t-2} \\ \text{type } B \text{ average action at time } t - 1 &= w_B \sum_{i=n/2+1}^n \frac{s_{M_i,t-1}}{n/2} + (1 - w_B) \theta_{t-2} \\ &\approx w_B \theta_{t-1} + (1 - w_B) \theta_{t-2}. \end{aligned}$$

Then, since $w_A > w_B$, the public knows two distinct linear combinations of θ_{t-1} and θ_{t-2} . The state θ_{t-1} is identified from these. So the member of the public can form a very precise estimate of θ_{t-1} —which, recall, is as well as she can hope to do. The key force is that the two groups of media outlets give different mixes of the old information and the more recent state, and by understanding this, the public can infer both. Indeed, to recover θ_{t-1} , the public puts a negative weight on the actions of media outlets of type B , which allows it to subtract off old information and focus on the recent state, θ_{t-1} . One can show that if, in contrast, agents are naive, *e.g.* if they think that all of the estimates of the media are uncorrelated (or only mildly correlated)

conditional on the state, they will put positive weights on their observations and will again be bounded from learning the state.

This illustration relied on a particular network with several special features: a very “central” source, one-directional links, and no communication among the media outlets or the public. We will show that the same considerations determine learning quality in a large class of random networks in which agents have many neighbours, with complex connections among them. Quite generally, if each neighbourhood contains a diversity of signal types, agents can concentrate on new developments in the state while filtering out old, less relevant information and thus estimate the changing state as accurately as physical constraints allow.

1.2. Outline

Section 2 sets up the basic model and discusses its interpretation. Section 3 defines equilibrium, shows its existence, and characterises it. Section 4 reports our main theoretical results on the quality of information aggregation. In Section 5, we discuss learning outcomes under a variety of non-Bayesian models. Section 6 defines and analyses social influence. Section 7 relates our model and results to the social learning literature. In Section 8, we describe our numerical exercise with network data from Indian villages and discuss a simple extension to multi-dimensional states to interpret our results on signal diversity.

2. MODEL

2.1. State of the world

There is a discrete set of instants, $\mathcal{T} = \mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$. At each time $t \in \mathcal{T}$, there is a *state*, a random variable θ_t taking values in \mathbb{R} . This state evolves as an AR(1) stochastic process. That is,

$$\theta_{t+1} = \rho\theta_t + v_{t+1}, \quad (2.1)$$

where ρ is a constant with $0 < |\rho| \leq 1$ and $v_{t+1} \sim \mathcal{N}(0, \sigma_v^2)$ are independent *innovations*. When $|\rho| < 1$ we have the explicit formula

$$\theta_t = \sum_{\ell=0}^{\infty} \rho^\ell v_{t-\ell},$$

and thus the state at any time t has the stationary distribution $\theta_t \sim \mathcal{N}(0, \sigma_v^2/(1 - \rho^2))$. We maintain the normalisation throughout that innovations have variance 1, *i.e.* $\sigma_v = 1$.

We will occasionally examine an alternative specification (making our departure from the main model explicit) where there is a starting time, so that $\mathcal{T} = \mathbb{Z}_{\geq 0} = \{0, 1, 2, \dots\}$, and the state process is defined as in (2.1) starting at time 0 with some specified distribution for θ_0 .

2.2. Information and observations

There is a set of *nodes* is $N = \{1, 2, \dots, n\}$. Each node i can be thought of as a location, and is associated with a set $N_i \subseteq N$ of nodes that i can observe, called its *neighbourhood*.⁶

Each node is populated by a sequence of *agents* in overlapping generations. For each time t , there is a node- i agent, labelled (i, t) , who takes that node’s time- t action $a_{i,t}$. This agent is

6. For all results, a node i ’s neighbourhood can, but need not, include i itself.

born at time $t - m$ and has m periods to observe the actions taken in her neighbourhood before she acts. Thus, when taking her action, the agent (i, t) knows $a_{j,t-\ell}$ for all nodes $j \in N_i$ and all lags $\ell \in \{1, 2, \dots, m\}$. We call m the *memory*; it reflects how many periods of actions in her neighbourhood an agent passively observes before acting. (See Figure 2 for an illustration.) One interpretation is that a node corresponds to a role in an organisation. A worker in that role has some time to observe colleagues in related roles before choosing a once-and-for-all action herself. Much of our analysis is done for an arbitrary finite m ; we view the restriction to finite memory as useful for avoiding technical complications, but because m can be arbitrarily large, this restriction has little substantive content.⁷

In addition to social information from her neighbourhood, each agent also observes a private signal,

$$s_{i,t} = \theta_t + \eta_{i,t},$$

where the error term $\eta_{i,t} \sim \mathcal{N}(0, \sigma_i^2)$ has a variance $\sigma_i^2 > 0$ that depends on the node but not on the time period. All the errors $\eta_{i,t}$ and state innovations v_t are independent of one another. An agent's information is a vector consisting of her private signal and all of her social observations. An important special case will be $m = 1$, where agents observe only one period of others' actions before acting themselves, so that the agent's information is $((a_{j,t-1})_{j \in N_i}, s_{i,t})$.

The *network* $G = (N, E)$ is the set of nodes N together with the (fixed) set of *links* E , defined as the subset of pairs $(i, j) \in N \times N$ such that $j \in N_i$. This network (also called a *graph*), which determines the observation structure, is common knowledge, as is the informational environment (*i.e.* the joint distribution of all exogenous random variables).

An *environment* is specified by (G, σ) , where $\sigma = (\sigma_i)_{i \in N}$ is the profile of signal variances.

2.3. Preferences and best responses

When an agent (i, t) makes her once-and-for-all choice $a_{i,t} \in \mathbb{R}$, her utility is given by

$$u_{i,t}(a_{i,t}) = -\mathbb{E}[(a_{i,t} - \theta_t)^2]. \tag{2.2}$$

7. It is worth noting that even when the memory m is small, observed actions can indirectly incorporate signals from much further in the past.




FIGURE 2

An illustration, with memory $m = 1$, of the overlapping generations at a node i . At time $t - 1$, agent (i, t) is born and observes contemporaneous actions in her neighbourhood. At time t , she observes her private signal $s_{i,t}$ and takes her action $a_{i,t}$

By a standard fact about squared-error loss functions, given the distribution of $(\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m$, the optimal choice of agent (i, t) is to set her action equal to her expectation of the state:

$$a_{i,t} = \mathbb{E}[\theta_t \mid \underbrace{(\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m}_{i\text{'s information}}, s_{i,t}]. \quad (2.3)$$

Here the notation $\mathbf{a}_{N_i,t'}$ refers to the vector $(a_{j,t'})_{j \in N_i}$ of time- t' actions in the agent's neighbourhood. An action can be interpreted as an agent's estimate of the state, and we will sometimes use this terminology.

The conditional expectation (2.3) depends, of course, on the prior of agent (i, t) about θ_t , which, under correctly specified beliefs, has distribution $\theta_t \sim \mathcal{N}(0, \sigma_v^2/(1 - \rho^2))$. We allow the prior to be any normal distribution or a diffuse improper prior.⁸ It saves on notation to analyse the case where all agents have improper priors. Because actions under a normal prior are related to actions under the improper prior by a simple linear bijection—and thus have the same information content for other agents—all results immediately extend to the general case.

The doubly infinite time axis introduces some subtleties into the definition of strategy profiles; complete details are formalised in [Supplementary Appendix OA3](#).

3. UPDATING AND EQUILIBRIUM

In this section, we study agents' learning behaviour and present a notion of stationary equilibrium. We begin with the canonical case of Bayesian agents with correct models of others' behaviour; we study other behavioural assumptions in Section 5 below.

3.1. Best-response behaviour

The first step is to analyse optimal updating behaviour in response to others' strategies. A strategy of an agent is *linear* if the action taken is a linear function of the variables she observes. We will analyse agents' best responses to linear strategies, showing that they are linear and computing them explicitly.⁹

Fix an agent (i, t) and some linear strategy profile played before time t . By (2.3), this agent's best-response action $a_{i,t}$ is her conditional expectation of θ_t given her information, $\mathbb{E}[\theta_t \mid (\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m, s_{i,t}]$. Each action before time t can be written as a (possibly infinite) sum of past signals $s_{j,t'}$. It follows that all random variables appearing in the conditional expectation are jointly Gaussian. That implies that $a_{i,t} = \mathbb{E}[\theta_t \mid (\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m, s_{i,t}]$ is an affine function of $s_{i,t}$ and $(\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m$ (see [Eaton, 1983](#), Section 4.3). We now analyse this conditional expectation in detail.

3.1.1. Actions as estimates of states: A key covariance matrix. Agents learn partly from past agents' actions, so the joint distribution of actions as estimates of the state at an arbitrary time t will be important to track. An arbitrary agent's scaled past action $\rho^\ell a_{i,t-\ell}$ gives an estimate of the state in the sense that $\mathbb{E}[\theta_t \mid \rho^\ell a_{i,t-\ell}] = \rho^\ell a_{i,t-\ell}$. Turning to second moments, define the

8. We take priors about θ_t , like the information structure and network, to be common knowledge.

9. This analysis applies both to the main $\mathcal{T} = \mathbb{Z}$ model and the alternative with $\mathcal{T} = \mathbb{Z}_{\geq 0}$. For a discussion of why it is natural to consider linear opponent strategies, see Section 3.2 below.

covariance matrix of the errors in these estimates over the most recent m periods:

$$\mathbf{V}_t = \text{Cov} \left((\rho^\ell a_{i,t-\ell} - \theta_t)_{\substack{i \in N \\ 0 \leq \ell \leq m-1}} \right).$$

In the case $m = 1$, we have $\mathbf{V}_t = \text{Cov}((a_{i,t} - \theta_t)_{i \in N})$. We will often refer to \mathbf{V}_t simply as the covariance matrix of the model, as it will play a central role in our subsequent analysis.

3.1.2. Best-response weights. The information of agent (i, t) at time t may be represented as a random vector

$$\mathbf{z}_{i,t} = \left((\rho^\ell a_{j,t-\ell})_{\substack{j \in N_i \\ 1 \leq \ell \leq m}}, s_{i,t} \right).$$

We will calculate the conditional expectation $\mathbb{E}[\theta_t | \mathbf{z}_{i,t}]$ in terms of a covariance matrix constructed from (i, t) 's observations,

$$\mathbf{C}_{i,t-1} = \text{Cov}(\mathbf{z}_{i,t} - \theta_t \mathbf{1}) = \begin{pmatrix} \rho^2 \mathbf{V}_{N_i,t-1} + \mathbf{1}\mathbf{1}^\top & \mathbf{0} \\ \mathbf{0} & \sigma_i^2 \end{pmatrix},$$

where $\mathbf{V}_{N_i,t-1}$ is the submatrix of \mathbf{V}_{t-1} corresponding to indices in i 's neighbourhood.¹⁰ Now we have the following formula for best-response actions, which, for simplicity, we give in the case where the agent has an improper prior¹¹ about θ_t :

$$a_{i,t} = \underbrace{\frac{\mathbf{1}^\top \mathbf{C}_{i,t-1}^{-1}}{\mathbf{1}^\top \mathbf{C}_{i,t-1}^{-1} \mathbf{1}}}_{\text{agent } (i,t)\text{'s weights}} \cdot \underbrace{\begin{pmatrix} \rho \mathbf{a}_{N_i,t-1} \\ \vdots \\ \rho^m \mathbf{a}_{N_i,t-m} \\ s_{i,t} \end{pmatrix}}_{\text{agent } (i,t)\text{'s observations}}. \quad (3.1)$$

Expression (3.1) is a linear combination of the agent's signal and the observed actions; the weights in this linear combination depend on the matrix \mathbf{V}_{t-1} , but *not* on realisations of any random variables. [Supplementary Appendix OA4.1](#) gives the details of the standard calculations underlying the formula.

We denote by $(\mathbf{W}_t, \mathbf{w}_t^s)$ a *weight profile* in period t , with $\mathbf{w}_t^s \in \mathbb{R}^n$ being the weights agents place on their private signals and \mathbf{W}_t being the weights they place on their other information.

3.1.3. The evolution of covariance matrices under best-response behaviour. Assuming agents best-respond according to the optimal weights just described in (3.1), we can compute the resulting next-period covariance matrix \mathbf{V}_t from the previous covariance matrix. Letting \mathcal{V} be the space of covariance matrices, this defines a map $\Phi : \mathcal{V} \rightarrow \mathcal{V}$, given by

$$\Phi : \mathbf{V}_{t-1} \mapsto \mathbf{V}_t. \quad (3.2)$$

10. We rewrite $\rho^\ell a_{j,t-\ell} - \theta_t = \rho(\rho^{\ell-1} a_{j,t-\ell} - \theta_{t-1}) - v_t$, where $\text{Var}[v_t] = 1$. The covariances of the term in parentheses are entries of \mathbf{V}_{t-1} . For the block structure, note the private signal errors $\eta_{i,t}$ are independent of events before t .

11. Our analysis extends immediately to any proper normal prior for θ_t : To get an agent's estimate of θ_t , the formula in (3.1) would simply be averaged with a constant term accounting for the prior, and everyone could invert this deterministic operation to recover the same information from others' actions.

This map gives the basic dynamics of the model: how an arbitrary variance–covariance matrix V_{t-1} maps to a new one when all agents best-respond to V_{t-1} . The variance–covariance matrix V_{t-1} (along with parameters of the model) determines (i) the weights agents place on their observations in (3.1), and (ii) the distributions of the random variables that are being combined in this expression. This yields the deterministic updating dynamic Φ . A consequence is that the weights agents place on observations are (commonly) known, and do not depend on any random realisations.

Example 1. We compute the map Φ explicitly in the case $m = 1$. We refer to the weight agent (i, t) optimally places on $\rho a_{j,t-1}$ as W_{ij} and the weight on $s_{i,t}$, her private signal, as w_i^s . Note we have, from (3.1) above, explicit expressions for these weights. Then

$$[\Phi(\mathbf{V})]_{ii} = (w_i^s)^2 \sigma_i^2 + \sum_{k,k'} W_{ik} W_{ik'} (\rho^2 V_{kk'} + 1) \quad \text{and} \quad [\Phi(\mathbf{V})]_{ij} = \sum_{k,k'} W_{ik} W_{jk'} (\rho^2 V_{kk'} + 1). \quad (3.3)$$

3.2. Stationary equilibrium in linear strategies

We will now turn our attention to *stationary equilibria in linear strategies*—ones in which all agents' strategies are linear with time-invariant coefficients—though, of course, we will allow all agents to consider deviating to arbitrary strategies, including nonlinear ones. Once we establish the existence of such equilibria, we will use the word *equilibrium* to refer to one of these unless otherwise noted.

A reason for focusing on equilibria in linear strategies comes from noting that, in the variant of the model with a starting time (*i.e.* the case $\mathcal{T} = \mathbb{Z}_{\geq 0}$) agents begin by using only private signals, and they do this linearly. After that, inductively applying the reasoning of Section 3.1, best-responses are linear at all future times. Taking time to extend infinitely backward is an idealisation that allows us to focus on exactly stationary behaviour.

We now show the existence of stationary equilibria in linear strategies.

Proposition 1. *A stationary equilibrium in linear strategies exists, and is associated with a covariance matrix $\hat{\mathbf{V}}$ such that $\Phi(\hat{\mathbf{V}}) = \hat{\mathbf{V}}$.*

The proof appears in [Appendix A](#), and we sketch the key ideas below.

At such an equilibrium, the covariance matrix \mathbf{V}_t and all agent strategies are time-invariant. Actions are linear combinations of observations with stationary weights, which we denote by \hat{W}_{ij} and \hat{w}_i^s . The form of these rules has some resemblance to static equilibrium notions studied in the rational expectations literature (*e.g.* [Vives, 1993](#); [Babus and Kondor, 2018](#); [Lambert *et al.*, 2018](#); [Mossel *et al.*, 2020](#)). It also has a similar form to the [DeGroot \(1974\)](#) and [Friedkin and Johnsen \(1997\)](#) updating rules, typically imposed as behavioural heuristics. In our dynamic environment, such a solution emerges as a stationary equilibrium.

3.2.1. Proof sketch for the existence result. The goal is to apply the Brouwer fixed-point theorem to show there is a covariance matrix $\hat{\mathbf{V}}$ that remains unchanged under updating. To find a convex, compact set to which we can apply the fixed-point theorem, we use the fact that when all agents best-respond to any beliefs about prior actions, all action variances lie in a compact set of positive numbers. This is because all agents' actions must be at least as precise in estimating θ_i as their private signals, and cannot be more precise than estimates given perfect knowledge of θ_{t-1} combined with the private signal. This establishes bounds on action variances. The Cauchy–Schwartz inequality then bounds covariances in terms of corresponding variances. All matrices

respecting these bounds constitute a compact, convex set containing the image of Φ . This and the continuity of Φ allow us to apply the Brouwer fixed-point theorem.

3.2.2. Other remarks. In the case of $m = 1$, we can use the formula of Example 1, equation (3.3), to write the fixed-point condition $\Phi(\hat{V}) = \hat{V}$ explicitly. More generally, for any m , equation (3.1) gives a formula in terms of \hat{V} for the weights \hat{W}_{ij} and \hat{w}_i^s in the best response to \hat{V} , and this can be used to describe the equilibrium \hat{V}_{ij} as solving a system of polynomial equations. These equations typically have large degree and cannot be solved analytically except in very simple cases, but they can readily be used to solve for equilibria numerically. A related feature of the model is that standard methods can easily be applied to estimate it and test hypotheses within it (see [Supplementary Appendix OA2](#) for details).

The main insight is that we can analyse equilibria through action covariances. This idea applies equally to many extensions and variations of our basic model, as illustrated by two examples: (1) We assume that agents observe neighbours perfectly, but one could define other observation structures. For instance, agents could observe actions with noise, or they could observe some set of linear combinations of neighbours' actions with noise. Similarly, agents could be observing predecessors' actions for heterogeneous durations before acting (*i.e.* node-specific m). (2) We assume agents are Bayesian and best-respond to the true distribution of actions, but the same proof would also show that equilibria exist under other behavioural rules (see Section 5.1).¹²

Proposition 1 shows that there exists a stationary linear equilibrium. We show later, as part of Proposition 2, that there is a *unique* stationary linear equilibrium in networks having a particular structure. In general, uniqueness of the equilibrium is an open question that we leave for future work.¹³ In Section 4.4 and [Supplementary Appendix OA5](#), we discuss the $\mathcal{T} = \mathbb{Z}_{\geq 0}$ variant of the model, which has a unique equilibrium, and relate it to our main model.

How much information does each agent need to play her equilibrium strategy? In a stationary equilibrium, she only needs to know the steady-state variance–covariance matrix \hat{V}_{N_i} in her neighbourhood. Then her problem of inferring θ_{t-1} becomes essentially a linear regression problem. If historical empirical data on neighbours' error variances and covariances are available, then \hat{V}_{N_i} can be estimated from such data.

4. HOW GOOD IS INFORMATION AGGREGATION IN EQUILIBRIUM?

In this section, we analyse the quality of information aggregation in stationary equilibrium.

First, recall that in any agent's time- t decision problem, θ_{t-1} is a sufficient statistic for social information, because the difference $\theta_t - \rho\theta_{t-1}$ is independent of all actions taken at or before time $t - 1$. Let the *social signal* of agent (i, t) be defined as her estimate of θ_{t-1} based on social information:

$$r_{i,t} = \mathbb{E}[\theta_{t-1} \mid (\mathbf{a}_{N_i,t-\ell})_{\ell=1}^m].$$

We will be interested in the error in this estimate:

12. What is important in the proof is that actions depend continuously on the covariance structure of an agent's observations; the action variances are uniformly bounded under the rule agents play; and there is a vanishing dependence of behaviour on the very distant past.

13. We have checked numerically that Φ is not, in general, a contraction in various norms (entrywise sup, Euclidean operator norm, etc.). In computing equilibria numerically for many examples, we have not been able to find a case of equilibrium multiplicity. Indeed, in all of our numerical examples, repeatedly applying Φ to an initial covariance matrix converges to the same fixed point for any starting conditions.

Definition 1. For a given strategy profile, define the aggregation error $\kappa_{i,t}^2 = \text{Var}(r_{i,t} - \theta_{t-1})$ to be the expected squared error in the social signal as a prediction of θ_{t-1} .

The aggregation error measures how well an agent can extract information from social observations. Note that agent i 's aggregation error is a monotone transformation of her expected utility.¹⁴

How efficient is aggregation? The environment features informational externalities: players do not internalise the impact of their learning rules on others' learning. Consequently, there is no reason to expect outcomes to be efficient in any exact sense. And we have seen that the details of equilibrium in a particular network can be complicated. However, it turns out that much more can be said about the behaviour of aggregation errors as neighbourhood sizes grow. In this section, we study the asymptotic efficiency of information aggregation. We give conditions under which aggregation error decays as quickly as physically possible, and different conditions under which it remains far from efficient levels even when agents have arbitrarily many observations. We discuss the case $m = 1$ for simplicity but the reasoning extends easily to other values of m .

4.1. A benchmark lower bound on aggregation error

A first observation is a lower bound on the aggregation error (in terms of an asymptotic rate as a function of a node's degree) under *any* behaviour of agents. This establishes a benchmark relative to which we can assess equilibrium outcomes.

Let d_i denote the out-degree of a node i .

Fact 1. Fix $\rho \in (-1, 1)$ as well as upper and lower bounds for private signal variances, so that $\sigma_i^2 \in [\underline{\sigma}^2, \bar{\sigma}^2]$ for all i . On any network and for all strategy profiles, we have $\kappa_{i,t}^2 \geq c/d_i$ for all i and t , where c is a constant that depends only on ρ , $\underline{\sigma}^2$, and $\bar{\sigma}^2$.

The lower bound is reminiscent of the central limit theorem: if an agent had d_i conditionally independent noisy signals about θ_{t-1} (e.g. by observing neighbours' private signals directly), then the variance of her estimate would be of order $1/d_i$. Fact 1 notes that it is not possible for aggregation errors to decay (as a function of degree) any faster than that.

For an intuition, imagine that an agent sees neighbours' private signals (not just actions) one period after they are received, and all other private signals two periods after they are received; this clearly gives an upper bound on the quality of the agent's possible aggregation given physical communication constraints. The information that is two periods old cannot be very informative about θ_{t-1} because of the movement in the state from period $t - 2$ to $t - 1$; a large constant number z of signals about θ_{t-1} would be better. Thus, a lower bound on aggregation error is given by the error that could be achieved with $d_i + z$ independent signals about θ_{t-1} of the best possible precision ($\underline{\sigma}^{-2}$). The bound follows from these observations.

4.2. Outline of results: When is aggregation comparable to the benchmark?

Fact 1 places a lower bound on aggregation error given the physical constraints. Even efficient learning could not do better than this bound. We examine when equilibrium learning can achieve aggregation of similar quality. More precisely, we ask when there is a stationary equilibrium where the aggregation error at node i satisfies $\hat{\kappa}_i^2 \leq C/d_i$ for all i , for some constant C .

14. In fact, for any decision dependent on θ_t , an agent is better off with a lower value of $\kappa_{i,t}^2$. This is a consequence of the fact that unidimensional Gaussian signals can be Blackwell ordered by their precision.

In Section 4.4, we establish a good-aggregation result: outcomes comparable to the benchmark are achieved in equilibrium in a class of networks. The key condition enabling the asymptotically efficient equilibrium outcome is called *signal diversity*: each individual has access to enough neighbours with multiple different kinds of private signals. The fact that neighbours use private information differently turns out to give the agents enough power to identify θ_{t-1} with equilibrium aggregation error that decays at a rate matching the lower bound of Fact 1 up to a multiplicative constant.

In Section 4.5, we turn to negative results. Without signal diversity, equilibrium aggregation can be extremely bad. Our first negative result shows that when signals are exchangeable, it may be that the aggregation error $\hat{\kappa}_i^2$ does not approach zero in any equilibrium, no matter how large neighbourhoods are, though a social planner could achieve good aggregation by prescribing different updating weights. We prove this in highly symmetric networks. Once we move away from such networks, one might ask whether diversity in individuals' network positions could play a role analogous to signal diversity and enable approximately efficient learning. Our next negative result shows that this is impossible. When signals are homogeneous and all agents' degrees in network G_n are bounded by $\bar{d}(n)$ (where $\bar{d}(n)$ is any unbounded sequence) then in any equilibrium, it cannot be that almost all aggregation errors are less than $C/\bar{d}(n)$ as the network grows, for any number $C > 0$ not depending on n .

4.3. Distributions of networks and signals

For our good-aggregation result, we study large populations and specify two aspects of the environment: *network distributions* and *signal distributions*. In terms of network distributions, we work with a standard type of random network model—a stochastic block model (see, e.g. Holland et al., 1983; Abbe, 2017). It makes the structure of equilibrium tractable while also allowing us to capture rich heterogeneity in network positions. We also specify *signal distributions*: how signal precisions are allocated to agents, in a way that may depend on network position. We now formalise these two primitives of the model and state the assumptions we work with.

Fix a set of *network types* $k \in \mathcal{K} = \{1, 2, \dots, K\}$. For each pair of network types, there is a given probability $p_{kk'}$ that each agent of network type k has a link to each agent of network type k' . An assumption we maintain on these probabilities is that each network type k observes at least one network type (possibly k itself) with positive probability. There is also a vector $(\alpha_1, \dots, \alpha_K)$ of *population shares* of each network type, which we assume are all positive. Jointly, $(p_{kk'})_{k,k' \in \mathcal{K}}$ and α specify the network distribution. These parameters can encode differences in expected degree and also features such as homophily (where some groups of types are linked to each other more densely than to others).

We next define signal distributions, which describe the allocation of signal variances to network types. Fix a finite set \mathbb{S} of private signal variances, which we call signal types.¹⁵ We let $q_{k\tau}$ be the share of agents of network type k with signal type τ ; then $(q_{k\tau})_{k \in \mathcal{K}, \tau \in \mathbb{S}}$ defines the signal distribution.

Let the nodes in network n be partitioned into the network types $N_n^1, N_n^2, \dots, N_n^K$, with the cardinality $|N_n^k|$ equal to $\lfloor \alpha_k n \rfloor$ or $\lceil \alpha_k n \rceil$ (rounding so that there are n agents in the network). We (deterministically) set the signal variances σ_i^2 equal to elements of \mathbb{S} in accordance with the signal shares (again rounding as needed). Let $(G_n)_{n=1}^\infty$ be a sequence of directed or undirected

15. The assumptions of finitely many signal types and network types are for technical convenience only, and could be relaxed.

random networks with these nodes, so that $i \in N_n^k$ and $j \in N_n^{k'}$ are linked with probability $p_{kk'}$; these link realisations are all independent.

In our setting, a *stochastic block model* D is specified by the linking probabilities $(p_{kk'})_{k,k' \in \mathcal{K}}$, the type shares α , and the signal distribution $(q_{k\tau})_{k \in \mathcal{K}, \tau \in \mathcal{S}}$. We let $(G_n(D), \sigma_n(D))$ denote the environment (*i.e.* the network and the signal variances) in a random realisation. We say that a network type k contains a signal type τ if $q_{k\tau} > 0$.

Definition 2. *A stochastic block model satisfies signal diversity if each network type has a positive probability of linking with at least one network type containing two distinct signal types.*

4.4. Good aggregation under diverse signals

Our first main result is that signal diversity is sufficient for good aggregation in the networks described in the previous section. Aggregation error decays at a rate C/d_i for each node i independently of the structural properties of the network.

We first define a notion of good aggregation for an agent in terms of a bound on that agent's aggregation error.

Definition 3. *Given $\varepsilon > 0$, we say that agent i achieves the ε -aggregation benchmark in a given equilibrium if the aggregation error satisfies $\hat{\kappa}_i^2 \leq \varepsilon$.*

We say an event (indexed by n) occurs *asymptotically almost surely* if the probability of the event converges to 1 as $n \rightarrow \infty$.

Theorem 1. *Fix any stochastic block model D satisfying signal diversity. There exists $C > 0$ such that asymptotically almost surely the environment $(G_n(D), \sigma_n(D))$ has an equilibrium where all agents achieve the C/n -aggregation benchmark.*

So for large enough n , societies with signal diversity are very likely to aggregate information very well. The uncertainty in this statement is over the network, as there is always a small probability of a realised network which prevents learning (*e.g.* an agent has no neighbours). We give an outline of the argument next, and the proof appears in [Appendix B](#).

The constant C in the theorem statement can depend on the stochastic block model D . However, given any compact set of stochastic block models D , we can choose a single $C > 0$ for which the result holds uniformly across D .¹⁶ Thus, the theorem can be applied without detailed information on how the random graphs are generated, as long as some bounds are known about which models are possible.

4.4.1. Discussion of the proof. To give intuition for [Theorem 1](#), we first describe why the theorem holds on the complete network¹⁷ with two signal types A and B in the $m = 1$ case. This echoes the intuition of the example in the Introduction. We then discuss the challenges involved in generalising the result to our general stochastic block model networks, and the techniques we use to overcome those challenges.

Consider a time- t agent, (i, t) . Recall that the social signal $r_{i,t}$ is the optimal estimate of θ_{t-1} based on the actions (i, t) has observed in her neighbourhood. In the complete network, all agents have the same social signal, which we call r_t .¹⁸

16. The reason is that the distribution of aggregation errors is upper hemicontinuous in model parameters, so if the desired bounds hold for each point in a compact set, they can be made uniform.

17. Note this is a special case of the stochastic block model.

18. In particular, agent (i, t) sees everyone's past action, including the one taken last period at the same node.

At any equilibrium, each agent’s action is a weighted average of her private signal and this social signal.

$$a_{i,t} = \hat{w}_i^s s_{i,t} + (1 - \hat{w}_i^s) r_t. \quad (4.1)$$

The weights on the two random variables on the right-hand side sum to 1 because both $s_{i,t}$ and r_t are unbiased estimates of θ_t , and so is the left-hand side $a_{i,t}$. The weight \hat{w}_i^s on the private signal depends on the precision of this signal relative to the social signal. We call the weights used by agents of the two distinct signal types \hat{w}_A^s and \hat{w}_B^s . Suppose signal type A is more precise than signal type B , so that $\hat{w}_A^s > \hat{w}_B^s$.

Now, turning our attention to the next period of updating, observe that each time- $(t + 1)$ agent can compute two averages of the time- t actions—one for each signal type. Using (4.1) to rewrite $a_{i,t}$ and then plugging in $s_{i,t} = \theta_t + \eta_{i,t}$:

$$\begin{aligned} \text{type } A \text{ average action at time } t &= \frac{1}{n_A} \sum_{i:\sigma_i^2=\sigma_A^2} a_{i,t} = \hat{w}_A^s \theta_t + (1 - \hat{w}_A^s) r_t + O_p(n^{-1/2}), \\ \text{type } B \text{ average action at time } t &= \frac{1}{n_B} \sum_{i:\sigma_i^2=\sigma_B^2} a_{i,t} = \hat{w}_B^s \theta_t + (1 - \hat{w}_B^s) r_t + O_p(n^{-1/2}). \end{aligned}$$

Here n_A and n_B denote the numbers of agents of each type (recalling we assumed each type is a positive share of the population size, n). The $O_p(n^{-1/2})$ error terms¹⁹ come from the average signal noises $\eta_{i,t}$ of agents in each group; the bound holds with high probability by the central limit theorem. In other words, each time- $(t + 1)$ agent can obtain precise estimates of two different convex combinations of θ_t and r_t . Because the two weights, \hat{w}_A^s and \hat{w}_B^s , are distinct, she can approximately (up to signal error) solve for θ_t as a linear combination of the average actions taken by each type she observes. It follows the agent must have an estimate at least as precise as what she can obtain by the strategy we have described, and will thus be very close the benchmark. Since the equilibrium in question was arbitrary, this shows that aggregation approaches the benchmark in any equilibrium. The estimator of θ_t in this strategy places negative weight on $(1/n_B) \sum_{i:\sigma_i^2=\sigma_B^2} a_{i,t}$, thus *anti-imitating* the agents of signal type B —those with the less precise private signal. The logic of Proposition 3 in Section 5.2 implies that anti-imitation necessarily occurs in any equilibrium in which agents aggregate information precisely.

To extend the ideas just presented to the more general setting of Theorem 1, we need to show that each individual observes a large number of neighbours of at least two signal types who also have similar social signals. More precisely, the proof shows that agents with the same network type have highly correlated social signals. Showing this is much more subtle than it was in the above illustration. In general, the social signals in an arbitrary network realisation are endogenous objects that depend to some extent on all the links.

A key insight allowing us to overcome this difficulty is a useful general fact about sufficiently dense stochastic block models: despite a lot of idiosyncratic randomness in direct connections, the law of large numbers implies the number of paths of length two between any agent i of type k and any agent j of type k' going through an agent of type k'' is nearly determined by the types k , k' , and k'' , with a small relative error.²⁰ We can leverage this to deduce some important facts

19. The notation means the errors are bounded by $Cn^{-1/2}$ for a $C > 0$ with high probability (Janson, 2011).

20. For simplicity, we first present the argument in a random graph family where the number of two-step paths is nearly deterministic. The argument extends to a larger class of models where the same property applies to longer paths, as we discuss in the next subsection.

about the updating map Φ (recall Section 3.1) in the realised random network, and specifically about the evolution of social signals.

In particular, if we look at the set of covariance matrices where all social signals are close to perfect, we can show that the composition $\Phi^2 := \Phi \circ \Phi$ maps this set to itself. In other words, if social signals are very precise, then they will remain very precise two periods later. If the two-step path counts were determined by types exactly, it would not be too difficult to show this by elaborating the reasoning in the complete graph example, because neighbours of the same type would be effectively exchangeable. We show that despite the path counts being known only approximately, the desired conclusion holds. This is nontrivial because the weights agents use in their updating—and thus the evolution of social signals—could depend sensitively on realised network structure; small relative errors could matter. A key step is to develop results on matrix perturbations to show that small relative changes in the network actually do not affect Φ^2 too much. A fixed-point theorem then implies there is a fixed point of Φ^2 in the set of outcomes with very precise social signals. With some further analysis we can deduce that this implies the existence of an equilibrium (corresponding to a fixed point of Φ) with nearly perfect aggregation.

4.4.2. Sparser random graphs. In the random graphs, we have defined in Section 4.3, the group-level linking probabilities ($p_{kk'}$) are, for simplicity, held fixed as n grows. This yields expected degrees that grow linearly in the population size, which may not be the desired asymptotic model. We can, however, establish versions of our results in a class of models much more flexible with respect to degrees. While it is important to have neighbourhoods “large enough” (*i.e.* growing in n) to permit the application of laws of large numbers, their rate of growth can be considerably slower than linear. For example, our proof can be extended directly to degrees that scale as n^α for any $\alpha > 0$ to show that asymptotically almost surely, there exists an equilibrium where the C/n^α -aggregation benchmark is achieved for all agents. Instead of studying Φ^2 and two-step paths, one can extend the same sort of analysis to the L -fold composition Φ^L , which reflects L -step paths. In order to do this, one uses the fact that for L larger than $1/\alpha$, the number of paths of length L between any two nodes is determined by the types involved in the path with a small relative error. Elaborating the proof of the theorem above, we can then characterise the behaviour of Φ^L and finally deduce the claimed aggregation property for Φ .

4.4.3. The good-aggregation outcome as a unique prediction. The theorem above says good aggregation is supported in an equilibrium but does not state that this is the unique equilibrium outcome. To deal with this issue, we study the alternative model with $\mathcal{T} = \mathbb{Z}_{\geq 0}$ (where agents begin with only their own signals and then best-respond to the previous distribution of behaviour at each time). We show that, as $n \rightarrow \infty$, the long-run outcomes of this model get arbitrarily close to the good-aggregation stationary equilibrium of Theorem 1 under the same conditions. Thus, even if there were other equilibria of the stationary model, they could not be approached via the natural iterative procedure coming from the $\mathcal{T} = \mathbb{Z}_{\geq 0}$ model. Formal statements and details are in [Supplementary Appendix OA5](#).

4.5. Aggregation under homogeneous signals

Having established conditions for good aggregation under signal diversity, we now explore what happens without signal diversity. Our general message is that aggregation is worse.

To gain an intuition for this, note that it is essential to the argument described in the previous subsection that different agents have different signal precisions. Recall the complete network case. From the perspective of an agent $(i, t + 1)$, the fact that type A and type B neighbours place different weights on the social signal r_t keeps their behaviour from being collinear, and

allows (i, t) to separate θ_t from a confound. In that example, if type A and B agents had the same signal types, they would use the same weights, and our agent trying to learn from them would face a collinearity problem.

We begin by studying graphs having a symmetric structure and show that learning outcomes are necessarily bounded very far from good aggregation. We then turn to arbitrary large graphs and prove a lower bound on aggregation error that implies the homogeneous signals regime has, quite generally, worse outcomes for some agents than those achieved by everyone in our good-aggregation result.

4.5.1. Aggregation in networks with symmetric neighbours.

Definition 4. *A network G has symmetric neighbours if, whenever $j, j' \in N_i$ for some i , then $N_j = N_{j'}$.*

In the undirected case, the graphs with symmetric neighbours are the complete network and complete bipartite networks.²¹ For directed graphs, the condition allows a larger variety of networks.

Proposition 2. *Consider a sequence $(G_n)_{n=1}^{\infty}$ of strongly connected graphs with symmetric neighbours. Assume that all signal variances are equal, and that $m = 1$. Then there is a unique equilibrium on each G_n , and there exists an $\varepsilon > 0$ such that the ε -aggregation benchmark is not achieved by any agent i at this equilibrium for any n .*

All agents have nonvanishing aggregation errors at the unique equilibrium. So all agents learn poorly compared with the diverse signals case. The proof of this proposition, and the proofs of all subsequent results, appear in [Supplementary Appendix OA4](#).

This failure of good aggregation is not due simply to a lack of sufficient information in the environment: On the complete graph with exchangeable (*i.e.* nondiverse) signals, a social planner who set weights for all agents could achieve ε -aggregation for any $\varepsilon > 0$ when n is large. See [Supplementary Appendix OA7](#) for a formal statement, proof and numerical results.²² In this sense, the social learning externalities are quite severe: a small change in weights for each individual could yield a very large benefit in a world of homogeneous signal types.

We now give intuition for Proposition 2. In a graph with symmetric neighbours and homogeneous signals, in the unique equilibrium,²³ actions of any agent's neighbours are exchangeable. So Bayesian estimates (and thus actions) must weight all neighbours equally, which prevents the sort of inference of the most recent state illustrated in Section 4.4. This is easiest to see on the complete graph, where *all* observations are exchangeable. So, in any equilibrium, each agent's action at time t is equal to a weighted average of her own signal and the average action $(1/|N_i|) \sum_{j \in N_i} a_{j,t-1}$:

$$a_{i,t} = \hat{w}_i^s s_{i,t} + (1 - \hat{w}_i^s) \frac{1}{|N_i|} \sum_{j \in N_i} a_{j,t-1}. \quad (4.2)$$

By iteratively using this equation, we can see that actions must place substantial weight on the average of signals from, *e.g.* two periods ago, and indeed further back. Note that all signals $s_{j,t'}$ at past times t' take the form $\theta_{t'} + \eta_{i,t'}$. Thus, although the effect of *signal errors* $\eta_{i,t'}$ vanishes

21. These are both special cases of our stochastic block model from Section 4.4, so Theorem 1 applies to these network structures when signal diversity is satisfied.

22. We thank Alireza Tahbaz-Salehi for suggesting this analysis.

23. The proof of the proposition establishes uniqueness by showing that Φ is a contraction in a suitable sense.

(by averaging) as n grows large, the correlated error from *past changes in the state* v_t never “washes out” of estimates, and this is what prevents vanishing aggregation errors.

The bad-aggregation result as stated applies to exactly homogeneous signal types only. In fact, in finite networks we need sufficiently heterogeneous signals to avoid bad learning outcomes; this is illustrated in [Supplementary Appendix OA1](#). In Section 4.6, we discuss the welfare implications of this failure of aggregation.

As a consequence of Theorem 1 and Proposition 2, we can give an example where making one node’s private information less precise helps all agents.

Corollary 1. *There exists a network G , a vector of signal precisions σ , and an agent $i \in G$ such that increasing σ_i^2 yields a Pareto improvement at the unique equilibrium.*

To prove the corollary, we consider the complete graph with homogeneous signals and large n . By Proposition 2, all agents have nonvanishing aggregation errors. If we instead give agent 1 a very uninformative signal, all players can anti-imitate agent 1 and achieve vanishing aggregation errors. When the signals at the initial configuration are sufficiently imprecise, this gives a Pareto improvement. There are also examples where severing links in the observational network can yield a Pareto improvement, as reported in an earlier version of the present paper ([Dasaratha et al., 2018](#)).

4.5.2. Aggregation in arbitrary networks. Section 4.5 showed aggregation errors are nonvanishing when signal endowments and neighbourhoods are symmetric. A natural question is whether asymmetry in network positions can substitute for asymmetry in signal endowments. In Section 4.4, the key point was that different neighbours’ actions were informative about different linear combinations of θ_i and older information, and this permitted filtering. Perhaps different network positions can achieve the same effect?

We thus move to *arbitrary* networks and show a weaker but much more general result. Consider any sequence of equilibria on any networks with symmetric signal endowments. Our result here is that no equilibrium achieves C/n -aggregation for almost all agents, no matter what C is. In particular, this implies that the rate of learning (as n grows) is slower than at the good-learning equilibrium with diversity of signal endowments from Theorem 1. Moreover, if degrees are bounded above by some $\bar{d}(n)$ growing at rate slower than n , we prove the stronger statement that no equilibrium achieves $C/\bar{d}(n)$ -aggregation for almost all agents.

Theorem 2. *Let $C > 0$. Let $(G_n)_{n=1}^\infty$ be an arbitrary sequence of networks and suppose all private signals have variance σ^2 . If all agents’ in-degrees and out-degrees are bounded above by some $\bar{d}(n) \rightarrow \infty$, then in any sequence of equilibria, the aggregation error $\hat{\kappa}_i^2$ is greater than $C/\bar{d}(n)$ for a nonvanishing fraction of agents i .*

In addition to considering arbitrary networks, we allow the memory m to be an arbitrary positive integer. Because the assumptions are much weaker, we obtain a weaker conclusion than in Proposition 2. While Proposition 2 shows that aggregation errors are nonvanishing, this theorem shows that aggregation errors cannot vanish quickly, but does not rule out aggregation errors vanishing more slowly.

The basic intuition is that to avoid putting substantial weight on θ_{t-2} , an agent at time t must anti-imitate some neighbours. If all or almost all neighbours achieve C/n -aggregation for some C and have identical types of private signals, there is not much diversity among neighbours. So more and more anti-imitation is needed as n grows large in the sense that the total positive weight and total negative weight on neighbours both grow large. But then the contribution to the agent’s variance from neighbours’ private signal errors cannot vanish quickly.

We can combine Theorems 1 and 2 to compare the value of signal diversity and network diversity. With diversity of signal endowments, there exists a $C > 0$ such that asymptotically almost surely there is a good-learning equilibrium achieving the C/n -aggregation benchmark for all agents under the stochastic block model. With exchangeable signals, it is not possible to find equilibria achieving the same aggregation rate in n under *any* sequence of networks. Thus, Theorem 2 shows that network heterogeneity cannot improve learning outcomes as much as signal heterogeneity. Section 8.1 complements the asymptotic results with numerical results in finite networks. It shows that in our model on real-world (highly asymmetric) social networks, signal heterogeneity improves learning outcomes much more than choosing a very favourable network structure but homogeneous signals.

4.6. *The welfare loss associated with homogeneity*

The results derived so far in this section show that there is a qualitative difference in how well agents are able to infer recent states across the homogeneous and heterogeneous signal settings. How important is this difference for welfare? We illustrate next that the welfare loss associated with signal homogeneity can be arbitrarily severe.

To gain an intuition for this, note that with homogeneous signals, period- t actions are confounded by previous states. These confounds include θ_{t-2} , which all $t - 1$ agents use in the same way (as illustrated in the example of the Introduction). But the confounds also include θ_{t-3} , which could not be filtered out by $t - 1$ agents, and so forth. The more weight agents place on social information (*i.e.* the more informative the past is), the more severe this confounding is. If the state is highly persistent and private signals are not very precise, then the confounds from periods even very long ago are substantial. The following corollary quantifies this effect.

Corollary 2. *Consider a complete graph with all signal variances equal to σ^2 , and let $m = 1$. Then, in any symmetric strategy profile,*

$$\text{Var}(a_{i,t} - \theta_t) \geq \frac{(1 - \hat{w}^s)^2}{1 - (1 - \hat{w}^s)^2 \rho^2},$$

where \hat{w}^s is the weight agents place on their own signals. As $\rho \rightarrow 1$ from below and $\sigma^{-2} \rightarrow 0$, agent i 's action error in the unique equilibrium tends to infinity. Moreover, this convergence is uniform in n .²⁴

The corollary guarantees that we can choose (σ^{-2}, ρ) so that the error is arbitrarily large, uniformly in n . In contrast, recall that our main positive result shows that the C/n -aggregation benchmark would be achieved with signal heterogeneity.²⁵ When this benchmark is achieved, each individual obtains a variance $\text{Var}(a_{i,t} - \theta_t)$ that is at worst 1 if n is large enough.²⁶ This bound on variance does not depend on σ^2 or ρ . Thus *welfare can be arbitrarily worse in environments with signal homogeneity compared to ones with heterogeneity.*

In large complete graphs with homogeneous signals, we can explicitly characterise the limit action variance (and therefore welfare). Let V^∞ denote the limit, as n grows large, of $\text{Var}(a_{i,t} - \theta_t)$. Let Cov^∞ denote the limit covariance of any two agents' errors. By direct

24. For any \underline{v} , there are $\rho < 1$ and $\bar{\sigma}^{-2} > 0$ such that if $\rho > \underline{\rho}$ and $\sigma^{-2} < \bar{\sigma}^{-2}$, then $\text{Var}(a_{i,t} - \theta_t) \geq \underline{v}$ for all n .

25. For example, by making half the agents' signals strictly worse.

26. Note the agent can use the estimate of last period's state, which has an error of order C/n . If the agent simply set her action equal to this estimate, then she would achieve $\text{Var}(\theta_t - \theta_{t-1}) = 1$, since the state innovation has variance 1. Additionally using her private signal does strictly better than this.

computation using equation (3.3), these can be seen to be related by the following equations, which have a unique solution:

$$V^\infty = \frac{1}{\sigma^{-2} + (\rho^2 \text{Cov}^\infty + 1)^{-1}}, \quad \text{Cov}^\infty = \frac{(\rho^2 \text{Cov}^\infty + 1)^{-1}}{[\sigma^{-2} + (\rho^2 \text{Cov}^\infty + 1)^{-1}]^2}. \quad (4.3)$$

These equations also let us extend Corollary 2 beyond the complete graph. The V^∞ and Cov^∞ solving (4.3) describe the limits of all variances and covariances in any graph with symmetric neighbours where degrees tend uniformly to infinity.²⁷ As $\sigma^{-2} \rightarrow 0$ and $\rho \rightarrow 1$ from below, equations (4.3) show that V^∞ and therefore Cov^∞ diverge to infinity, just as in the complete-network case. This shows the welfare loss from homogeneity can also be arbitrarily severe in graphs with symmetric neighbours and large degrees.

5. THE IMPORTANCE OF UNDERSTANDING CORRELATIONS

In the positive result on achieving the C/n -aggregation benchmark (Theorem 1), a key aspect of the argument involved agents filtering out confounding information from their neighbours' estimates—*i.e.* responding in a sophisticated way to the correlation structure of those estimates. In this section, we demonstrate that this sort of behaviour is essential for nearly perfect aggregation, and that more naively imitative heuristics yield outcomes far from the benchmark. Empirical studies have found evidence (depending on the setting and the subjects) consistent with both equilibrium behaviour and naive inference in the presence of correlated observations (*e.g.* Eyster *et al.*, 2015; Enke and Zimmermann, 2019; Dasaratha and He, 2021).

We begin with a canonical model of agents who do not account for correlations among their neighbours' estimates conditional on the state, and show by example that naive agents achieve much worse learning than Bayesian agents, and thus have nonvanishing aggregation errors. We then formalise the idea that accounting for correlations in neighbours' actions is crucial to reaching the benchmark. This is done by demonstrating a general lack of good aggregation by agents who use imitative strategies, rather than filtering in a sophisticated way. Finally, we show that even in fixed, finite networks, any positive weights chosen by optimising agents will be Pareto-dominated.

5.1. Naive agents

In this part, we introduce agents who misunderstand the distribution of the signals they are facing and who therefore do not update as Bayesians with a correct understanding of their environment. We consider a particular form of misspecification that simplifies solving for equilibria analytically.²⁸

Definition 5. *We call an agent naive if she believes that all neighbours choose actions equal to their private signals and maximises her expected utility given these incorrect beliefs.*

Equivalently, a naive agent believes her neighbours all have empty neighbourhoods. This is the analogue, in our model, of “best-response trailing naive inference” (Eyster and Rabin, 2010).

27. Indeed, it can be deduced (as in the proof of Corollary 2) that agents' actions are equal to an appropriately discounted sum of past $\theta_{t-\ell}$, up to error terms (arising from $\eta_{i,t-\ell}$) that vanish asymptotically. The weights on past states are the same as in the complete-network case, which is why the characterisation of (4.3) applies.

28. There are a number of possible variants of our behavioural assumption, and it is straightforward to numerically study alternative specifications of behaviour in our model (Alatas *et al.*, 2016 consider one such variant).




FIGURE 3

Bayesian and naive learning on a complete graph and $n = 600$ agents divided into two groups of equal size. The plot shows the aggregation error in group A as group B's private signal variance varies, fixing group A's private signal variance at $\sigma_A^2 = 2$

So naive agents understand that their neighbours' actions from the previous period are estimates of θ_{t-1} , but they think these are conditionally independent given the state, and that the precision of each estimate is equal to the signal precision of the corresponding agent. They then play their expectation of the state given this misspecified theory of others' play.

In Figure 3, we compare Bayesian and naive learning outcomes. We consider a complete network with 600 agents and $\rho = 0.9$. Half of agents have signal variance $\sigma_A^2 = 2$, while we vary the signal variance σ_B^2 of the remaining agents. The figure shows the average social signal variance for the group of agents with private signal variance $\sigma_A^2 = 2$. It suggests that naive agents learn substantially worse than rational agents, whether signals are diverse or not. We prove this holds for general stochastic block models and provide formulas for variances under naive learning in [Supplementary Appendix OA6](#).

5.2. More general learning rules: Understanding correlation is essential for good aggregation

We now show more generally that a sophisticated response to correlation is needed to achieve vanishing aggregation errors on any sequence of growing networks. To this end, we make the following definition:

Definition 6. *The steady state associated with weights \mathbf{W} and \mathbf{w}^s is the (unique) covariance matrix \mathbf{V}^* such that if actions have a variance-covariance matrix given by $\mathbf{V}_t = \mathbf{V}^*$ and next-period actions are set using weights $(\mathbf{W}, \mathbf{w}^s)$, then $\mathbf{V}_{t+1} = \mathbf{V}^*$ as well.*

In this definition of steady state, instead of best-responding to others' actual distributions of play, agents use exogenous weights \mathbf{W} in all periods.

By a straightforward application of the contraction mapping theorem, if agents use any non-negative weights under which covariances remain bounded at all times, there is a unique steady state.

Consider a sequence of networks $(G_n)_{n=1}^\infty$ with n agents in G_n .

Proposition 3. *Fix any sequence $(V^*(n))_{n=1}^\infty$, with each $V^*(n)$ being a steady state under non-negative weights in the network G_n . Suppose that all private signal variances are bounded below by $\underline{\sigma}^2 > 0$ and that all agents place weight at most $\bar{w} < 1$ on their private signals. Then there is an $\varepsilon > 0$ such that, for all n , the ε -aggregation benchmark is not achieved by any agent i at the steady state $V^*(n)$.*

The essential idea is that at time $t + 1$, observed time- t actions all put weight on actions from period $t - 1$, which causes θ_{t-1} to have a (positive weight) contribution to all observed actions. Agents do not know θ_{t-1} and, with positive weights, cannot take any linear combination that would recover it. Even with a very large number of observations, this confound prevents agents from learning the time- t state precisely.

We now explain why we impose an assumption of all weights on private signals being bounded away from 1. If there were many autarkic agents who simply reported their private signals (*i.e.* placed weight 1 on these signals), some other agent could learn well without adjusting for correlations by observing the autarkic agents. Note that in this case, all of the autarkic agents would have nonvanishing aggregation errors. This illustrates that a weaker conclusion than that of the proposition can be established more generally. If we did not impose a bound on private signal weights, learning would fail in the weaker sense that *some* agent must fail to achieve the ε -aggregation benchmark for small enough ε .

On undirected networks, the proposition implies that aggregation errors do not vanish under naive inference or under various other specifications of non-Bayesian inference implying nonnegative weights. Moreover, the same argument shows that in any sequence of *Bayesian* equilibria on undirected networks where all agents use positive weights, no agent can learn well.

5.3. Without anti-imitation, outcomes are Pareto-inefficient

The previous section argued that anti-imitation is critical to achieving vanishing aggregation errors. We now show that even in small networks, where that benchmark is not relevant, any equilibrium without anti-imitation is Pareto-inefficient relative to another steady state. This result complements our asymptotic analysis by showing a different sense (relevant for small networks) in which anti-imitation is necessary to make the best use of information.

Proposition 4. *Suppose the network G is strongly connected and some agent has more than one neighbour. Given any naive equilibrium or any Bayesian equilibrium where all weights are positive, the action variances at that equilibrium are Pareto-dominated by action variances at another steady state.*

The basic argument behind Proposition 4 is that if agents place marginally more weight on their private signals, this introduces more independent information that eventually benefits everyone. In a review of sequential learning experiments, [Weizsäcker \(2010\)](#) finds that subjects weight their private signals more heavily than is optimal (given the empirical behaviour of others they observe). Proposition 4 implies that in our environment with optimising agents, it is actually welfare-improving for individuals to “overweight” their own information relative to best-response behaviour.

The condition on equilibrium weights says that no agent anti-imitates any of her neighbours. This assumption makes the analysis tractable, but we believe the basic force also works in finite networks with some anti-imitation. In the proof in [Supplementary Appendix OA4](#), we state and prove a more general result where weights are nonnegative but need not all arise from Bayesian or naive updating.

5.3.1. Proof sketch. The idea of the proof for the Bayesian equilibrium case is to begin at the steady state and then marginally shift each agent’s weights toward her private signal. This means agents’ actions are less correlated but, by the envelope theorem, not significantly worse in the next period. We show that if all agents continue using these new weights, the decreased correlation eventually benefits everyone. To do this, we use the absence of anti-imitation, which implies a certain monotonicity in the updating function whereby the initial decrease in correlation results in all agents’ variances decreasing.

The proof in the naive case is simpler. Here a naive agent is overconfident about the quality of her social information, so she would benefit from shifting some weight from her social information to her signal. This deviation also reduces her correlation with other agents, so it is Pareto-improving.

6. SOCIAL INFLUENCE

A canonical question about learning in networks is how much influence various agents have in affecting aggregate behaviour. This is a focus of studies including [DeMarzo *et al.* \(2003\)](#) and [Golub and Jackson \(2010\)](#) in the DeGroot model with an unchanging state. In this section, we define a suitable analogue of social influence for our dynamic environment. We then study how an agent’s influence depends on her signal precision and degree. We find that, relative to benchmark results from the DeGroot model, influence is more sensitive to signal precisions, while social connectedness plays a similar role in both models.

6.1. Defining social influence

We define the *total influence* of node i in a stationary equilibrium with weights (\hat{W}, \hat{w}^s) to be the total weight that all actions place on the private signal of agent (i, t) . The total influence measures the total increase in actions if $s_{i,t}$ increases by 1 (due to an idiosyncratic shock).²⁹ At equilibrium, the total influence of i is:

$$\text{TI}(i) = \sum_{j \in N} \sum_{k=0}^{\infty} (\rho^k \hat{W}^k)_{ji} \hat{w}_i^s.$$

This expression for total influence is a version of Katz–Bonacich centrality with respect to the matrix \hat{W} of weights. The decay parameter is the persistence ρ of the of the AR(1) state process.

We define the *social influence* of i to be the total weight that all actions *in future periods* place on the private signal of agent (i, t) . At equilibrium, the social influence of i is:

$$\text{SI}(i) = \sum_{j \in N} \sum_{k=1}^{\infty} (\rho^k \hat{W}^k)_{ji} \hat{w}_i^s = \text{TI}(i) - \hat{w}_i^s.$$

The social influence measures the influence of an agent at node i on other agents. Social influence and total influence differ only by the weight (i, t) places on her own current signal, because an agent’s signal realisation does not affect others’ actions in the same period. Note that agent i ’s social influence depends on the weight \hat{w}_i^s she places on her own signal as well as the weights agents place on each others’ actions.

29. Note that in a stationary equilibrium, this depends on the node and not the time, so we speak interchangeably of the influence of a node and that of an agent at this node.

The next result, which follows from Proposition 1 on equilibrium existence, shows that the summation that defines social influence is guaranteed to converge at equilibrium, which makes social influence (and similarly total influence) well defined.³⁰

Proposition 5. *The social influence $SI(i)$ is well defined at any equilibrium and is equal to $[\mathbf{1}^\top (\mathbf{I} - \rho \hat{\mathbf{W}})^{-1} - \mathbf{1}^\top]_i \hat{w}_i^s$.*

We show this as follows: if social influence did not converge, some agents would have actions with very large variances (because their actions would depend sensitively on small idiosyncratic shocks). But then these agents would have simple deviations that would improve their accuracy, such as following their private signals. So this could not happen in equilibrium. Once the infinite series defining social influence is shown to be convergent, the proposition follows by a standard Neumann series identity.

In general, social influence can be negative: an agent's net effect on others can be in the opposite direction of her signal.

6.2. Which agents are influential?

We now ask how the social influence $SI(i)$ of an agent depends on her signal precision and degree. To facilitate the most direct comparison with standard results in models with a fixed state, such as DeMarzo *et al.* (2003), we focus on cases where social influences are positive.

To examine the effect of signal precision on social influence, we first study complete networks with $n \geq 2$ agents and two private signal variances: half the agents have more precise signals, and the other half have less precise signals. We call the two groups' signal variances σ_A^2 and σ_B^2 and the corresponding agents' social influences $SI(A)$ and $SI(B)$. We show that the ratio between the two groups' social influences in equilibrium is larger than the ratio between their signal precisions (whenever the imprecise group has positive social influence).

Proposition 6. *On a complete network with $m = 1$ and signal variances $\sigma_A^2 < \sigma_B^2$, in the unique equilibrium it holds that*

$$\frac{SI(A)}{SI(B)} > \frac{\sigma_A^{-2}}{\sigma_B^{-2}}$$

whenever $SI(B) > 0$.

The proposition says that increasing a group's precision increases their influence more than proportionately. As we have seen in our main results, if the precision difference is large enough, then it is optimal to place zero or negative weight on the less precise group. The result says that even before this happens, imprecision reduces a group's influence considerably—and, as we will discuss below, more than in benchmark models of social influence.

The proposition assumes the network is complete, but numerical evidence suggests that on other networks, too, agents with more precise signals tend to be much more influential. We simulate a configuration model with $n = 40$ nodes, each with degree $d = 5$.³¹ Nodes are randomly assigned to have a precise signal with variance σ_A^2 or an imprecise signal with variance σ_B^2 (with equal probability).

30. Since $\hat{\mathbf{W}}$ can contain both positive and negative numbers, some of them potentially large, it is not immediately obvious that the summation converges.

31. This model works by creating n nodes, each with d "stubs" sticking out of it, and then performing a random matching of the stubs to create a graph. See Jackson (2010), Section 4.5.10, for details.

TABLE 1

The table shows how far influence ratios are from a benchmark of being proportional to precision

Precision	σ_B^2									
	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1	2.14	4.53	9.06	168.09	-29.69	-15.54	-8.22	-7.41	-6.19
1		1	1.80	3.15	5.62	9.95	26.35	-29.89	-15.23	-13.67
1.5			1	1.64	2.56	3.80	6.27	10.64	44.97	-31.32
2				1	1.51	2.22	3.13	4.35	7.05	11.34
2.5					1	1.43	2.04	2.69	3.88	5.50
σ_A^2 3						1	1.38	1.86	2.46	3.35
3.5							1	1.34	1.74	2.27
4								1	1.30	1.67
4.5									1	1.28
5										1

Notes: We use a configuration model with a regular network and heterogeneous signal variances; there are $n = 40$ agents and the degree is $d = 5$. Agents are randomly assigned to signal variances σ_A^2 or σ_B^2 . Each entry is computed from 100 runs with persistence $\rho = 0.9$. Each table entry reports the ratio $R_\sigma = (SI(A)/SI(B))/(\sigma_A^{-2}/\sigma_B^{-2})$ for the precision parameters corresponding to that entry.

We are interested in the ratio $SI(A)/SI(B)$ in this more complicated environment. If social influence were approximately proportional to precision, then $SI(A)/SI(B)$ would be approximately $\sigma_A^{-2}/\sigma_B^{-2}$. To assess by how much the influence of the precise group exceeds the level suggested by this benchmark, we will look at the ratio

$$R_\sigma = \frac{SI(A)/SI(B)}{\sigma_A^{-2}/\sigma_B^{-2}}.$$

Table 1 reports this ratio over 100 runs of the simulation model for various pairs of σ_A^2 and σ_B^2 , each in the interval $[0.5, 5]$. The entries of the table would be equal to 1 if influence is proportional to precision. Instead, all off-diagonal entries are greater than one (or negative), meaning social influence depends more (and often much more) on signal precision than in the proportional benchmark.

Having examined how influence depends on precisions, we turn to how it depends on degrees. We again use a configuration model, which allows us to fix any desired empirical degree distribution and generate the graphs uniformly conditional on the degrees. We will find that social influence depends less on degree than on precision. We simulate a configuration model with $n = 40$ nodes, each randomly assigned degrees d_A or d_B (with equal probability of each) and with $\sigma^2 = 2$ for all agents. Table 2 reports the ratio

$$R_d = \frac{SI(A)/SI(B)}{d_A/d_B}.$$

over 100 runs of the simulation model for degrees between 1 and 10. Again, the entries would be equal to 1 if social influence were proportional to degree. Social influence is indeed approximately proportional to degree: the entries in the table range between 0.91 and 1.11.

Remark 1. A simple intuition explains why social influence depends more on private information than on network position. Increasing an agent’s private signal precision and her degree both tend to make her action more accurate. Increasing private signal precision has the additional effect of increasing an agent’s weight on her private information, which is recent and independent of other agents’ actions. This provides more reason for others to place weight on

her actions, amplifying the effect of the increased accuracy. In contrast, increasing degree tends to make an agent place more weight on her social information, which is older and more correlated with others. This countervails the effect of increased accuracy, making the agent a less appealing source for others.

The exercises so far varied only one of signal precision or degree, and we now explore how social influence depends on precision and degree jointly. To do so, we compute equilibrium social influences on 5000 networks with $n = 40$ agents in each. Each agent is randomly assigned a degree chosen uniformly from $\{1, 2, \dots, 7\}$ and a private signal variance chosen uniformly and independently from $\{0.5, 1, \dots, 3.5\}$. Networks are then drawn via the configuration model. Figure 4 plots the level curves for average social influence conditional on node attributes. The steepness of the level curves shows that social influence again depends more on signal variance than degree, especially when signals are less precise.

6.3. Comparison with a DeGroot benchmark

The results above are interesting to compare with those of canonical network models with a fixed state. A relevant benchmark is a version of the DeGroot model studied by DeMarzo *et al.* (2003). Agents start with an improper prior, receive independent normal private signals s_i (with different precisions) about the state once, and then each takes an action $a_{i,0}$ equal to her expectation of the state θ . After this, agents observe their neighbours' actions and take actions $a_{i,1}$, which are Bayesian expectations of the state θ given their observations. In all subsequent periods $t > 1$, agents observe their neighbours' actions $a_{j,t-1}$ and take actions $a_{i,t}$ as if $a_{j,t-1}$ had the same distribution as j 's private signal. That is, they naively repeat their optimal strategy from the first period, which DeMarzo *et al.* (2003) interpret as a quasi-Bayesian, boundedly rational procedure.

One natural measure of social influence is the influence of s_i on the long-run consensus estimate $\lim_{t \rightarrow \infty} a_{j,t}$ held by any agent. In an undirected, connected, and aperiodic network, this limit exists and the influence of agent i is proportional to her private signal precision σ_i^{-2} and to her degree d_i . Compared to this benchmark, social influence in our changing-state model is more sensitive to signal precision (in the complete graph and in our simulations for configuration

TABLE 2
The table shows how far influence ratios are from a benchmark of being proportional to degree

		d_B									
Degree		1	2	3	4	5	6	7	8	9	10
d_A	1	1	1.11	1.02	0.96	0.94	0.92	0.96	1.00	1.03	1.09
	2		1	1.02	1.01	0.98	0.96	0.92	0.93	0.91	0.95
	3			1	1.01	1.01	1.01	0.98	0.96	0.94	0.93
	4				1	1.01	1.01	1.00	0.99	0.98	0.96
	5					1	1.01	1.01	1.02	1.01	1.00
	6						1	1.01	1.01	1.01	1.01
	7							1	1.01	1.01	1.02
	8								1	1.01	1.01
	9									1	1.01
	10										1

Notes: We use a configuration model with two possible degrees on $n = 40$ agents with homogeneous signal variance $\sigma^2 = 2$. Agents are randomly assigned to degrees d_A or d_B . Each entry is computed from 100 runs with persistence $\rho = 0.9$. Each table entry reports the ratio $R_d = (\text{SI}(A)/\text{SI}(B))/(d_A/d_B)$ for the degree parameters corresponding to that entry.




FIGURE 4

Level curves for average social influence of agents in a configuration model with 5000 networks with $n = 40$ agents in each and persistence $\rho = 0.9$. Degrees are chosen uniformly from $\{1, 2, \dots, 7\}$ and private signal variances are chosen uniformly from $\{0.5, 1, \dots, 3.5\}$. The figure shows level curves for average social influence (drawn via cubic interpolation) on a log–log plot. If proportional changes in degree and signal variances mattered equally, these level curves would have slope 1.

models). On the other hand, the dependence of social influence on degree is very similar to the DeGroot benchmark—approximately proportional. To summarise, influence depends more on an agent’s private information, while the dependence on network position is remarkably similar. The difference between the benchmark and our model is explained in Remark 1.

7. RELATED LITERATURE

Whether decentralised communication can facilitate efficient adaptation to a changing world is a fundamental question in economic theory, related to questions raised by Hayek (1945)³² and central to certain applied problems, *e.g.* in real business cycle models with consumers and firms learning about evolving states.³³ Nevertheless, there is relatively little modelling of Bayesian learning of dynamic states in the large literature on social learning and information aggregation in networks, whose most relevant papers we now review.³⁴

32. “If . . . the economic problem of society is mainly one of rapid adaptation to changes in the particular circumstances of time and place . . . there still remains the problem of communicating to [each individual] such further information as he needs.” Hayek’s main concern was aggregation of information through markets, but the same questions apply more generally.

33. See Angeletos and La’O (2010) for a survey of related models that are used to study real business cycles. More recent developments include Angeletos and Lian (2018) and Molavi (2019), with the latter allowing a form of misspecification.

34. For more complete surveys of different parts of this literature, see, among others, Acemoglu and Ozdaglar (2011), Golub and Sadler (2016), and Mossel and Tamuz (2017). See Moscarini *et al.* (1998) for an early model in a binary-action environment, where it is shown that a changing state can break information cascades.

Play in the stationary linear equilibria of our model closely resembles behaviour in the DeGroot (1974) model, where agents update by linearly aggregating network neighbours' past estimates, with constant weights on neighbours over time. DeMarzo *et al.* (2003), in a Gaussian environment with an unchanging state, derive DeGroot learning as the Bayesian behaviour in the first round of communication, and use that as a foundation for a DeGroot rule as a boundedly rational heuristic. Molavi *et al.* (2018) present new bounded-rationality foundations for the DeGroot rule. Our different environment offers a different foundation for averaging rules with time-invariant weights: as a stationary equilibrium of a stationary environment.³⁵ Though the updating rule resembles those studied in fixed-state environments, we have stressed that the learning implications are quite different.

Several recent papers in engineering and computer science study dynamic environments similar to ours. Shahrampour *et al.* (2013) study an exogenous-weights version, interpreted as a set of Kalman filters under the control of a planner. They bound measures of welfare in terms of the persistence of the state process (ρ) and network properties, such as the spectral gap. Frongillo *et al.* (2011) study a $\rho = 1$ model of the state. They characterise the steady-state distribution of behaviour for any weights, and calculate equilibrium weights on a complete network, which they show are inefficient. Our Proposition 4 documents a related inefficiency; the quality of equilibrium learning in large, incomplete networks and social influence in equilibrium are topics not considered in these papers. In economics, Alatas *et al.* (2016) perform an empirical exercise in a similar model with a quasi-Bayesian learning rule. Their estimation assumes agents ignore the correlations between social observations, similarly to our naive models.³⁶ Our results show that the degree of rationality can be pivotal for the outcomes of such processes, and provide foundations for structural inference to test various behavioural assumptions.

Our results about when agents learn well are related to two phenomena that have played an important role in the social learning literature. One theme in this literature is that heterogeneity—in agents' neighbourhoods or preferences—can be helpful for learning. A manifestation of this is the usefulness of *sacrificial lambs* (typically studied in sequential social learning models with a fixed state): a small set of agents who observe nobody can help everyone else learn well, because their actions are then informative only about their private signals, and unconfounded by an information cascade (SgROI, 2002; Arieli and Mueller-Frank, 2019). Heterogeneity in preferences can serve a similar purpose: if preferences have full support, there is a positive probability that preference bias counteracts available social information, causing an agent to follow her private signal (Goeree *et al.*, 2006; Lobel and Sadler, 2016). A crucial difference is that our mechanism does not rely on any agents simply revealing their private signals: heterogeneity helps by changing how neighbours use their *social* information, which in turn aids an agent in inferring a common confound.³⁷

Second, a robust aspect of rational learning in sequential models is the phenomenon of anti-imitation, as discussed, *e.g.* by Eyster and Rabin (2014). They give general conditions for fully Bayesian agents to anti-imitate in a standard sequential model. We find that anti-imitation is

35. Indeed, agents behaving according to the DeGroot heuristic in other environments might have to do with their experiences in stationary environments where it is closer to optimal.

36. The paper's focus is estimating parameters of social learning rules using data from Indonesian villages, where agents are trying to estimate each other's wealth.

37. A bit farther afield, in Sethi and Yildiz (2012), learning outcomes when two individuals repeatedly learn from each other depend on whether their (heterogeneous) priors are independent or correlated; the common thread is that a natural assumption about agents' attributes (independent priors in their case) leads to an identification problem. The mechanics are otherwise quite different.

also an important feature in our dynamic model, and in our context is crucial for good learning. Despite this similarity, there is an important contrast between our environment and standard sequential models. In those models, while rational agents *do* prefer to anti-imitate, individuals and society as a whole can often obtain good outcomes using heuristics without any anti-imitation: for instance, by combining one’s own private signal with the information that can be inferred from a single neighbour. [Acemoglu *et al.* \(2011\)](#) and [Lobel and Sadler \(2015\)](#) show that such a heuristic leads to asymptotic learning in a sequential model. Our dynamic learning environment is different, as shown in Proposition 3: to have any hope of approaching good aggregation benchmarks, agents must respond in a sophisticated way, with anti-imitation, to their neighbours’ (correlated) estimates.

8. DISCUSSION AND EXTENSIONS

8.1. *Aggregation and its absence without asymptotics: Numerical results*

The message of Section 4 is that signal diversity enables good aggregation, and signal homogeneity obstructs it. The theoretical results were asymptotic, and relied on various assumptions about network structure. It is natural to ask whether our main conclusions hold up in realistic finite networks. To analyse this, we numerically study equilibria of our model on graphs reflecting social relationships measured in Indian villages ([Banerjee *et al.*, 2013](#)). This subsection briefly summarises our findings; we describe the exercise fully in [Supplementary Appendix OA1](#).

We examine the benefits of signal heterogeneity for equilibrium aggregation. The network data are essentially the only empirical input to our exercise.³⁸ Given a network, we compute equilibria using our model and parameters chosen for illustration. We compare two environments that differ in signal allocations: (i) a homogeneous case, with all signal variances set to 2, and (ii) a heterogeneous case, where half of the nodes have a signal variance greater than 2 (which we vary) and half of the nodes have a signal variance less than 2.³⁹

We first compare the value of a good network with the value of heterogeneous signals. Some networks have better learning than others even with homogeneous signals. We define the *network-driven variation* in learning to be the standard deviation of learning quality (aggregation error) across villages in the homogeneous case. Our main finding is that increasing the private signal variance for half of the agents by 50%, and reducing the signal variance of the others to keep total information constant, changes social signal error variance by 6.5 times the network-driven variation. In fact, introducing this amount of private signal heterogeneity improves learning much more than the most favourable network among the villages.

Though the asymptotic prediction changes starkly depending on whether signal precisions are identical or not, considerable heterogeneity is actually required to achieve the benefits of signal diversity in a finite network. Starting from homogeneous signals and increasing signal diversity, aggregation error changes very slightly at first. Once the variance of the less precise signal has increased by 50% relative to the starting point, learning quality has moved about halfway to what is achievable with the most extreme signal heterogeneity.

38. In particular, we have no data on signal qualities; when we introduce signal heterogeneity, we simply posit that households without electricity have worse access to external information.

39. We choose the larger signal variance so that the average precision in each village is $\frac{1}{2}$, which holds the total inflow of information constant in a sense made precise in the appendix.

8.2. *Multidimensional states and informational specialisation*

Our formal analysis assumed a one-dimensional state and one-dimensional signals, which varied only in their precisions. Our message about the value of diversity is, however, better interpreted in a mathematically equivalent multidimensional model.

Consider Bayesian agents who learn and communicate about two independent dimensions simultaneously, each one working as in our model. If all agents have equally precise signals about both dimensions, then society may not learn well about either of them. In contrast, if half the agents have superior signals about one dimension and inferior signals about the other (and the other half has the reverse), then society can learn well about both dimensions. Thus, the designer has a strong preference for an organisation with informational specialisation where some, but not all, agents are expert in a particular dimension.⁴⁰

Of course, there are many familiar reasons for specialisation in having precise information about an issue. For instance, it may be that specialisation is technologically efficient, or makes it easier to provide incentives. Crucially, specialisation is valuable in our setting for a distinct reason: it helps agents with their inference problems.

More generally, one could readily extend our model and equilibrium concept to a multi-dimensional state $\theta_t \in \mathbb{R}^d$ and arbitrary Gaussian signals about it, with flexible correlations. We would expect to find suitable generalisations of the basic message that sufficient diversity within neighbourhoods (in terms of signal types) facilitates learning. The assumption that agents know neighbours' signal distributions is clearly very helpful for tractability; it would be interesting to consider models in which agents are also uncertain about these distributions.

APPENDIX A. EXISTENCE OF EQUILIBRIUM: PROOF OF PROPOSITION 1

Recall from Section 3.1 the map Φ , which gives the next-period covariance matrix $\Phi(V_t)$ for any V_t . The expression given there for this map ensures that its entries are continuous functions of the entries of V_t . Our strategy is to show that this function maps a convex, compact set, \mathcal{K} , to itself, which, by Brouwer's fixed-point theorem, ensures that Φ has a fixed point \hat{V} . We will then argue that this fixed point corresponds to a stationary linear equilibrium.

We begin by defining the compact set \mathcal{K} . Recalling Section 3.1, entries of V_t are covariances between pairs of neighbour errors from any periods $t - \ell$ where $1 \leq \ell \leq m$. Let k, l be two indices of such actions, corresponding to actions taken at nodes i and j , respectively (at potentially different times), and let $\bar{\sigma}_i^2 = \max\{\sigma_i^2, \rho^{m-1}\sigma_i^2 + (1 - \rho^{m-1})/(1 - \rho)\}$. Now let $\mathcal{K} \subset \mathcal{V}$ be the subset of symmetric positive semi-definite matrices V_t such that, for any such k, l ,

$$V_{kk,t} \in \left[\min \left\{ \frac{1}{1 + \sigma_i^{-2}}, \frac{\rho^{m-1}}{1 + \sigma_i^{-2}} + \frac{1 - \rho^{m-1}}{1 - \rho} \right\}, \max \left\{ \sigma_i^2, \rho^{m-1}\sigma_i^2 + \frac{1 - \rho^{m-1}}{1 - \rho} \right\} \right]$$

$$V_{kl,t} \in [-\bar{\sigma}_i \bar{\sigma}_j, \bar{\sigma}_i \bar{\sigma}_j].$$

This set is closed and convex, and we claim that $\Phi(\mathcal{K}) \subset \mathcal{K}$.

To show this claim, we will first find upper and lower bounds on the variance of any neighbour's action (at any period in memory). For the upper bound, note that a Bayesian agent will not choose an action with a larger variance than her signal, which has variance σ_i^2 . For a lower

40. This raises important questions about what information agents would acquire, and whom they would choose to observe, which are the focus of a growing literature. For recent papers on this in the context of networks, see [Sethi and Yildiz \(2016\)](#) and [Myatt and Wallace \(2017\)](#), among others.

bound, note that if she knew the previous period's state and her own signal, then the variance of her action would be $1/(1 + \sigma_i^{-2})$. Thus an agent observing only noisy estimates of θ_t and her own signal can do no better.

By the same reasoning applied to the node- i agent from m periods ago, the error variance of $\rho^m a_{i,t-m} - \theta_t$ is at most $\rho^m \sigma_i^2 + (1 - \rho^m)/(1 - \rho)$ and at least $\rho^m/(1 + \sigma_i^{-2}) + (1 - \rho^m)/(1 - \rho)$. This establishes bounds on $V_{kk,t}$ for observations k from either the most recent or the oldest available period. The corresponding bounds from the periods between $t - m + 1$ and t are always weaker than at least one of the two bounds we have described, so we need only take minima and maxima over two terms.

This established the claimed bound on the variances. The bounds on covariances follow from Cauchy–Schwartz.

We have now established that there is a variance–covariance matrix \hat{V} such that $\Phi(\hat{V}) = \hat{V}$. By definition of Φ , this means there exists some weight profile (\hat{W}, \hat{w}^s) such that, when applied to prior actions that have variance–covariance matrix \hat{V} , produce variance–covariance matrix \hat{V} . However, it still remains to show that this is the variance–covariance matrix reached when agents have been using the weights (\hat{W}, \hat{w}^s) forever.

To show this, first observe that if agents have been using the weights (\hat{W}, \hat{w}^s) forever, the variance–covariance matrix V_t in any period is uniquely determined and does not depend on t ; call this \check{V} .⁴¹ This is because actions can be expressed as linear combinations of private signals with coefficients depending only on the weights. Second, it follows from our construction above of the matrix \hat{V} and the weights (\hat{W}, \hat{w}^s) that there is a distribution of actions where the variance–covariance matrix is \hat{V} in every period and agents are using weights (\hat{W}, \hat{w}^s) in every period. Combining the two statements shows that in fact $\check{V} = \hat{V}$, and this completes the proof. Note that this argument also establishes that the response profile we have constructed is a strategy profile: under the responses used, we can write formally the dependence of actions on all prior signals, and verify using the observations on decay of dependence across time that the formula is summable and hence defines unique actions.

APPENDIX B. PROOF OF THEOREM 1

B.1. Notation and key notions

Let \mathbb{S} be the (by assumption finite) set of all possible signal variances, and let $\bar{\sigma}^2$ be the largest of them. The proof will focus on the covariances of errors in social signals. Suppose that all agents have at least one neighbour. Take two arbitrary agents i and j . Recall that both $r_{i,t}$ and $r_{j,t}$ have mean θ_{t-1} , because each is an unbiased estimate⁴² of θ_{t-1} ; we will thus focus on the errors $r_{i,t} - \theta_{t-1}$. Let A_t denote the variance–covariance matrix $(\text{Cov}(r_{i,t} - \theta_{t-1}, r_{j,t} - \theta_{t-1}))_{i,j}$ and let \mathcal{W} be the set of such covariance matrices. For all i, j note that $\text{Cov}(r_{i,t} - \theta_{t-1}, r_{j,t} - \theta_{t-1}) \in [-\bar{\sigma}^2, \bar{\sigma}^2]$ using the Cauchy–Schwarz inequality and the fact that $\text{Var}(r_{i,t} - \theta_{t-1}) \in [0, \bar{\sigma}^2]$ for all i . This fact about variances says that no social signal is worse than putting all weight on an agent who follows only her private signal. Thus the best-response map Φ is well defined and induces a map $\check{\Phi}$ on \mathcal{W} .

Next, for any $\psi, \zeta > 0$ we will define the subset $\mathcal{W}_{\psi,\zeta} \subset \mathcal{W}$ to be the set of covariance matrices in \mathcal{W} such that both of the following hold:

41. The variance–covariance matrices are well defined because the (W, w^s) weights yield unambiguous strategy profiles in the sense of [Supplementary Appendix OA3](#).

42. This is because it is a linear combination, with coefficients summing to 1, of unbiased estimates of θ_{t-1} .

1. for any pair of distinct agents⁴³ $i \in G_n^k$ and $j \in G_n^{k'}$,

$$\text{Cov}(r_{i,t} - \theta_{t-1}, r_{j,t} - \theta_{t-1}) = \psi_{kk'} + \zeta_{ij}$$

where (i) $\psi_{kk'}$ depends only on the network types of the two agents (k and k' , which may be the same); (ii) $|\psi_{kk'}| < \psi$; and (iii) $|\zeta_{ij}| < \zeta$;

2. for any single agent $i \in G_n^k$,

$$\text{Var}(r_{i,t} - \theta_{t-1}) = \psi_k + \zeta_{ii}$$

where (i) ψ_k only depends on the network type of the agent; (ii) $|\psi_k| < \psi$, and (iii) $|\zeta_{ii}| < \zeta$.

This is the space of covariance matrices such that each covariance is split into two parts. Considering (1) first, $\psi_{kk'}$ is an effect that depends only on i 's and j 's network types, while ζ_{ij} adjusts for the individual-level heterogeneity arising from different link realisations. The description of the decomposition in (2) is analogous.

B.2. Proof strategy

A set $\mathcal{W}_{\bar{\psi}, \bar{\zeta}}$ of outcomes with good learning. Our goal is to show that as n grows large, there is an equilibrium in which $\text{Var}(r_{i,t} - \theta_{t-1})$ becomes very small, which then implies that the agents asymptotically learn. To this end we define a set of covariances with this property as well as some other useful properties. We will take $\bar{\psi}$ and $\bar{\zeta}$ to be arbitrarily small numbers and show that for large enough n , with high probability (which we abbreviate ‘‘asymptotically almost surely’’ or ‘‘a.a.s.’’) there is an equilibrium with a social error covariance matrix A_t in the set $\mathcal{W}_{\bar{\psi}, \bar{\zeta}}$. That will imply that, in this equilibrium, $\text{Var}(r_{i,t} - \theta_{t-1})$ becomes arbitrarily small as we take the constants $\bar{\psi}$ and $\bar{\zeta}$ to be small. In our constructions, the ζ_{ij} (resp., ζ_i) terms will be set to much smaller values than the $\psi_{kk'}$ (resp., ψ_k) terms, because group-level covariances are more predictable and less sensitive to idiosyncratic realisations than individual-level covariances.

Approach to showing that $\mathcal{W}_{\bar{\psi}, \bar{\zeta}}$ contains an equilibrium. To show that there is (a.a.s.) an equilibrium outcome with a social error covariance matrix A_t in the set $\mathcal{W}_{\bar{\psi}, \bar{\zeta}}$, the plan is to construct a set so that (a.a.s.) $\bar{\mathcal{W}} \subset \mathcal{W}_{\bar{\psi}, \bar{\zeta}}$ and $\tilde{\Phi}(\bar{\mathcal{W}}) \subset \bar{\mathcal{W}}$. This set will contain an equilibrium by the Brouwer fixed-point theorem, and therefore so will $\mathcal{W}_{\bar{\psi}, \bar{\zeta}}$.

To construct the set $\bar{\mathcal{W}}$, we will fix a positive constant β (to be determined later), and define

$$\bar{\mathcal{W}} = \mathcal{W}_{\beta/n, 1/n} \cup \tilde{\Phi}(\mathcal{W}_{\beta/n, 1/n}).$$

We will then prove that, for large enough n , (i) $\tilde{\Phi}(\bar{\mathcal{W}}) \subseteq \bar{\mathcal{W}}$ and (ii) for another suitable positive constant λ ,

$$\bar{\mathcal{W}} \subset \mathcal{W}_{\beta/n, \lambda/n}.$$

This will allow us to establish that (a.a.s.) $\bar{\mathcal{W}} \subset \mathcal{W}_{\bar{\psi}, \bar{\zeta}}$ and $\tilde{\Phi}(\bar{\mathcal{W}}) \subset \bar{\mathcal{W}}$, with $\bar{\psi}$ and $\bar{\zeta}$ being arbitrarily small numbers.

The following two lemmas will allow us to deduce (immediately after stating them) properties (i) and (ii) of $\bar{\mathcal{W}}$.

43. Throughout this proof, we abuse terminology by referring to agents and nodes interchangeably when the relevant t is clear or specified nearby.

Lemma B.1. *There is a function $\underline{\lambda}(\beta) \geq 1$ such that the following holds. For all large enough β and all $\lambda \geq \underline{\lambda}(\beta)$, for n sufficiently large we have $\tilde{\Phi}(\mathcal{W}_{\beta/n, 1/n}) \subset \mathcal{W}_{\beta/n, \lambda/n}$ with probability at least $1 - 1/n$.*

Lemma B.2. *For all large enough β , for n sufficiently large, $\tilde{\Phi}^2(\mathcal{W}_{\beta/n, 1/n}) \subset \mathcal{W}_{\beta/n, 1/n}$, with probability at least $1 - 1/n$.⁴⁴*

Putting these lemmas together, a.a.s. we have,

$$\tilde{\Phi}^2(\mathcal{W}_{\beta/n, 1/n}) \subset \mathcal{W}_{\beta/n, 1/n} \quad \text{and} \quad \tilde{\Phi}(\mathcal{W}_{\beta/n, 1/n}) \subset \mathcal{W}_{\beta/n, \lambda/n}.$$

From this it follows that $\bar{\mathcal{W}} = \mathcal{W}_{\beta/n, 1/n} \cup \tilde{\Phi}(\mathcal{W}_{\beta/n, 1/n})$ is mapped to a subset of itself by $\tilde{\Phi}$, and contained in $\mathcal{W}_{\beta/n, \lambda/n}$, as claimed.

Proving the lemmas by analysing how $\tilde{\Phi}$ and $\tilde{\Phi}^2$ act on sets $\mathcal{W}_{\psi, \zeta}$. The lemmas are about how $\tilde{\Phi}$ and $\tilde{\Phi}^2$ act on the covariance matrix \mathbf{A}_t , assuming it is in a certain set $\mathcal{W}_{\psi, \zeta}$, to yield new covariance matrices. Thus, we will prove these lemmas by studying two periods of updating. The analysis will come in five steps.

Step 1: No-large-deviations (NLD) networks and the high-probability event. Step 1 concerns the “with high probability” part of the lemmas. In the entire argument, we condition on the event of a *no-large-deviations (NLD)* network realisation, which says that certain realised statistics in the network (e.g. number of paths between two nodes) are close to their expectations. The expectations in question depend only on agents’ types. Therefore, on the NLD realisation, the realised statistics do not vary much based on which exact agents we focus on, but rather depend only on their types. Step 1 defines the NLD event E formally and shows that it has high probability. We use the structure of the NLD event throughout our subsequent steps, as we mention below.

Step 2: Weights in one step of updating are well-behaved. We are interested in $\tilde{\Phi}$ and $\tilde{\Phi}^2$, which describe how the covariance matrix \mathbf{A}_t of social signal errors changes under updating. How this works is determined by the “basic” updating map Φ , and so we begin by studying the weights involved in it and then make deductions about the implications for the evolution of the variance–covariance matrix \mathbf{A}_t .

The present step establishes that in one step of updating, the weight $W_{ij, t+1}$ that agent $(i, t+1)$ places on the action of another agent j in period t , does not depend too much on the identities of i and j . It only depends on their (network and signal) types. This is established by using our explicit formula for weights in terms of covariances. We rely on (i) the fact that covariances are assumed to start out in a suitable $\mathcal{W}_{\psi, \zeta}$, and (ii) our conditioning on the NLD event E . The NLD event is designed so that the network quantities that go into determining the weights depend only on the types of i and j (because the NLD event forbids too much variation within type). The restriction to $\mathbf{A}_t \in \mathcal{W}_{\psi, \zeta}$ ensures that covariances in the initial period t do not vary too much with type, either.

Step 3: Lemma B.1: $\tilde{\Phi}(\mathcal{W}_{\beta/n, 1/n}) \subset \mathcal{W}_{\beta/n, \lambda/n}$. Once we have analysed one step of updating, it is natural to consider the implications for the covariance matrix. Because we now have a bound on how much weights can vary after one step of updating, we can compute bounds on covariances. We show that if covariances \mathbf{A}_t are in $\mathcal{W}_{\beta/n, 1/n}$, then after one step, covariances

44. The notation $\tilde{\Phi}^2$ means the operator $\tilde{\Phi}$ applied twice.

are in $\mathcal{W}_{\beta/n,\lambda/n}$. Note that the introduction of another parameter λ on the right-hand side implies that this step might worsen our control on covariances somewhat, but in a bounded way.

Step 4: Weights in two steps of updating are well behaved. Here we establish that the statement made in Step 2 remains true when we replace $t + 1$ by $t + 2$. By the same sort of reasoning as in Step 2, an additional period of updating cannot create too much further idiosyncratic variation in weights. Proving this requires analysing the covariance matrices of various social signals (*i.e.* the \mathbf{A}_{t+1} that the updating induces), which is why we needed to do Step 3 first.

Step 5: Lemma B.2: $\tilde{\Phi}^2(\mathcal{W}_{\beta/n,1/n}) \subset \mathcal{W}_{\beta/n,1/n}$. Now we use our understanding of weights from the previous steps, along with additional structure, to show the key remaining fact. What we have established so far about weights allows us to control the weight that a given agent's estimate at time $t + 2$ places on the social signal of another agent at time t . This is Step 5(a). In the second part, Step 5(b), we use that to control the covariances in \mathbf{A}_{t+2} . It is important in this part of the proof that different agents have very similar "second-order neighbourhoods": the paths of length 2 beginning from an agent are very similar, in terms of their counts and what types of agents they go through. We use our control of second-order neighbourhoods, as well as the assumptions on variation across entries of \mathbf{A}_t to bound this variation well enough to conclude that $\mathbf{A}_{t+2} \in \mathcal{W}_{\beta/n,1/n}$.

B.3. Carrying out the steps

Step 1. Here we formally define the NLD event, which we call E . It is given by $E = \bigcap_{i=1}^5 E_i$, where the events E_i will be defined next.

(E_1) Let $X_{i,\tau k}^{(1)}$ be the number of agents having signal type τ and network type k who are observed by i . The event E_1 is that this quantity is close to its expected value in the following sense, simultaneously for all possible values of the subscript:

$$(1 - \zeta^2)\mathbb{E}[X_{i,\tau k}^{(1)}] \leq X_{i,\tau k}^{(1)} \leq (1 + \zeta^2)\mathbb{E}[X_{i,\tau k}^{(1)}].$$

(E_2) Let $X_{ii',\tau k}^{(2)}$ be the number of agents having signal type τ and network type k who are observed by *both* i and i' . The event E_2 is that this quantity is close to its expected value in the following sense, simultaneously for all possible values of the subscript:

$$(1 - \zeta^2)\mathbb{E}[X_{ii',\tau k}^{(2)}] \leq X_{ii',\tau k}^{(2)} \leq (1 + \zeta^2)\mathbb{E}[X_{ii',\tau k}^{(2)}].$$

(E_3) Let $X_{i,\tau k,j}^{(3)}$ be the number of agents having signal type τ and network type k who are observed by agent i and who observe agent j . The event E_3 is that this quantity is close to its expected value in the following sense, simultaneously for all possible values of the subscript:

$$(1 - \zeta^2)\mathbb{E}[X_{i,\tau k,j}^{(3)}] \leq X_{i,\tau k,j}^{(3)} \leq (1 + \zeta^2)\mathbb{E}[X_{i,\tau k,j}^{(3)}].$$

(E_4) Let $X_{ii',\tau k,j}^{(4)}$ be the number of agents having signal type τ and network type k who are observed by both agent i and i' and who observe j . The event E_4 is that this quantity is close to its expected value in the following sense, simultaneously for all possible values of the subscript:

$$(1 - \zeta^2)\mathbb{E}[X_{ii',\tau k',j}^{(4)}] \leq X_{ii',\tau k',j}^{(4)} \leq (1 + \zeta^2)\mathbb{E}[X_{ii',\tau k',j}^{(4)}].$$

(E_5) Let $X_{i,\tau k,jj'}^{(5)}$ be the number of agents of signal type τ and network type k who are observed by agent i and who observe both j and j' . The event E_5 is that this quantity is close to its expected value in the following sense, simultaneously for all possible values of the subscript:

$$(1 - \zeta^2)\mathbb{E}[X_{i,\tau k,jj'}^{(5)}] \leq X_{i,\tau k,jj'}^{(5)} \leq (1 + \zeta^2)\mathbb{E}[X_{i,\tau k,jj'}^{(5)}].$$

We claim that the probability of the complement of the event E vanishes exponentially. We can check this by showing that the probability of each of the E_i vanishes exponentially. For E_1 , for example, the bounds will hold unless at least one agent has degree outside the specified range. The probability of this is bounded above by the sum of the probabilities of each individual agent having degree outside the specified range. By Chebyshev's inequality, the probability a given agent has degree outside this range vanishes exponentially. Because there are n agents in G_n , this sum vanishes exponentially as well. The other cases are similar.

For the rest of the proof, we condition on the event E .

Step 2. As a shorthand, let $\psi = \beta/n$ for a sufficiently large constant β , and let $\zeta = 1/n$.

Lemma B.3. *Suppose that in period t the matrix $\mathbf{A} = \mathbf{A}_t$ of covariances of social signals satisfies $\mathbf{A} \in \mathcal{W}_{\psi,\zeta}$ and all agents are optimising in period $t + 1$. Then there is a γ so that for all n sufficiently large,*

$$\frac{W_{ij,t+1}}{W_{i'j',t+1}} \in \left[1 - \frac{\gamma}{n}, 1 + \frac{\gamma}{n}\right].$$

whenever i and i' have the same network and signal types and j and j' have the same network and signal types.

To prove this lemma, we will use the formula given by (3.1) for row i of the matrix \mathbf{W}_{t+1} :

$$W_{i,t+1} = \frac{\mathbf{1}^\top \mathbf{C}_{i,t}^{-1}}{\mathbf{1}^\top \mathbf{C}_{i,t}^{-1} \mathbf{1}}.$$

This says that in period $t + 1$, agent i 's weight on agent j is proportional to the sum of the entries of column j of $\mathbf{C}_{i,t}^{-1}$. We want to show that the change in weights is small as the covariances of observed social signals vary slightly. To do so we will use the Taylor expansion of $f(\mathbf{A}) = \mathbf{C}_{i,t}^{-1}$ around the covariance matrix $\mathbf{A}(0)$ at which all $\psi_{kk'} = 0$, $\psi_k = 0$ and $\zeta_{ij} = 0$.

We begin with the first partial derivative of f at $\mathbf{A}(0)$ in an arbitrary direction. Let $\mathbf{A}(x)$ be any perturbation of $\mathbf{A}(0)$ in one parameter, *i.e.* $\mathbf{A}(x) = \mathbf{A}(0) + x\mathbf{M}$ for some constant matrix \mathbf{M} with entries in $[-1, 1]$. Let $\mathbf{C}_i(x)$ be the matrix of covariances of the actions observed by i given that the covariances of agents' social signals were $\mathbf{A}(x)$. There exists a constant γ_1 depending only on the possible signal types such that each entry of $\mathbf{C}_i(x) - \mathbf{C}_i(x')$ has absolute value at most $\gamma_1(x - x')$ whenever both x and x' are small.

We will now show that the column sums of $\mathbf{C}_i(x)^{-1}$ are close to the column sums of $\mathbf{C}(0)_i^{-1}$. To do so, we will evaluate the formula

$$\frac{\partial f(\mathbf{A}(x))}{\partial x} = \frac{\partial \mathbf{C}_i(x)^{-1}}{\partial x} = \mathbf{C}_i(x)^{-1} \frac{\partial \mathbf{C}_i(x)}{\partial x} \mathbf{C}_i(x)^{-1} \quad (\text{B.1})$$

at zero. If we can bound each column sum of this expression (evaluated at zero) by a constant (depending only on the signal types and the number of network types K), then the first derivative of f will also be bounded by a constant.

Recall that \mathbb{S} is the set of signal types and let $S = |\mathbb{S}|$; index the signal types by numbers ranging from 1 to S . To bound the column sums of $\mathbf{C}_i(0)^{-1}$, suppose that the agent observes r_i agents from each signal type $1 \leq i \leq S$. Reordering so that all agents of each signal type are grouped together, we can write, for a suitable matrix \mathbf{Y} and vector \mathbf{z} :

$$\mathbf{C}_i(0) = \begin{pmatrix} Y_{11}\mathbf{1}_{r_1 \times r_1} + z_1 \mathbf{I}_{r_1} & Y_{12}\mathbf{1}_{r_1 \times r_2} & & Y_{S1}\mathbf{1}_{r_1 \times r_S} \\ Y_{12}\mathbf{1}_{r_2 \times r_1} & Y_{22}\mathbf{1}_{r_2 \times r_2} + z_2 \mathbf{I}_{r_2} & & \vdots \\ & & \ddots & \\ Y_{1S}\mathbf{1}_{r_S \times r_1} & \cdots & & Y_{SS}\mathbf{1}_{r_S \times r_S} + z_S \mathbf{I}_{r_S} \end{pmatrix}$$

Therefore, the covariance matrix $\mathbf{C}_i(0)$ can be written as a block matrix with blocks $Y_{\tau\tau'}\mathbf{1}_{r_\tau \times r_{\tau'}} + z_\tau \delta_{\tau\tau'} \mathbf{I}_{r_\tau}$ where $1 \leq \tau, \tau' \leq S$ and $\delta_{\tau\tau'} = 1$ for $\tau = \tau'$ and 0 otherwise.

We now have the following important approximation of the inverse of this matrix.⁴⁵

Lemma B.4. *Pinelis (2018)* Let \mathbf{C} be a block matrix with blocks given by

$$Y_{\tau\tau'}\mathbf{1}_{r_\tau \times r_{\tau'}} + z_\tau \delta_{\tau\tau'} \mathbf{I}_{r_\tau}$$

for all $\tau, \tau' \in \mathbb{S}$. As $n \rightarrow \infty$, the (τ, τ) block of \mathbf{C}^{-1} satisfies

$$\frac{1}{z_\tau} \mathbf{I}_{r_\tau} - \frac{1}{z_\tau r_\tau} \mathbf{1}_{r_\tau \times r_\tau} + O(1/n^2)$$

while the off-diagonal blocks are $O(1/n^2)$.

Proof. Note that the block (τ, τ') of \mathbf{C}^{-1} has the form

$$\mathbf{E}_{\tau\tau'}\mathbf{1}_{r_\tau \times r_{\tau'}} + d_\tau \delta_{\tau\tau'} \mathbf{I}_{r_\tau}$$

for some matrix \mathbf{E} and vector \mathbf{d} . Here δ denotes the Kronecker delta.

Therefore, the (τ, τ') block of $\mathbf{C}\mathbf{C}^{-1}$ can be written (using that $\mathbf{1}_{r \times r'}\mathbf{1}_{r' \times r''} = r'\mathbf{1}_{r \times r''}$) as

$$\begin{aligned} & \sum_{\hat{\tau}} (Y_{\tau\hat{\tau}}\mathbf{1}_{r_\tau \times r_{\hat{\tau}}} + z_\tau \delta_{\tau\hat{\tau}} \mathbf{I}_{r_\tau}) (\mathbf{E}_{\hat{\tau}\tau'}\mathbf{1}_{r_{\hat{\tau}} \times r_{\tau'}} + d_{\hat{\tau}} \delta_{\hat{\tau}\tau'} \mathbf{I}_{r_{\hat{\tau}}}) \\ &= \left(Y_{\tau\tau'} d_{\tau'} + \sum_{\hat{\tau}} (Y_{\tau\hat{\tau}} r_{\hat{\tau}} + \delta_{\tau\hat{\tau}} z_{\hat{\tau}}) \mathbf{E}_{\hat{\tau}\tau'} \right) \mathbf{1}_{r_\tau \times r_{\tau'}} + z_\tau d_\tau \delta_{\tau\tau'} \mathbf{I}_{r_\tau}. \end{aligned} \quad (\text{B.2})$$

For any vector $\mathbf{v} \in \mathbb{R}^{\mathbb{S}}$, let \mathbf{D}_v denote the diagonal matrix with v_τ in the (τ, τ) diagonal entry and $\mathbf{v} \circ \mathbf{v}'$ denote the pointwise product of two vectors. Breaking up the fact that (B.2) equals \mathbf{I} into its off-diagonal and diagonal parts, we have

$$\mathbf{Y}\mathbf{D}_d + (\mathbf{Y}\mathbf{D}_r + \mathbf{D}_z)\mathbf{E} = \mathbf{0} \text{ and } \mathbf{D}_d = \mathbf{D}_z^{-1}.$$

45. We are very grateful to Iosif Pinelis for this argument.

Hence,

$$\begin{aligned}
 E &= -(\mathbf{Y}\mathbf{D}_r + \mathbf{D}_z)^{-1}\mathbf{Y}\mathbf{D}_d \\
 &= -(\mathbf{I}_S + \mathbf{D}_r^{-1}\mathbf{Y}^{-1}\mathbf{D}_z)^{-1}(\mathbf{Y}\mathbf{D}_r)^{-1}\mathbf{Y}\mathbf{D}_z^{-1} \\
 &= -(\mathbf{I}_S + \mathbf{D}_r^{-1}\mathbf{Y}^{-1}\mathbf{D}_z)^{-1}\mathbf{D}_{z\circ r}^{-1} \\
 &= -\mathbf{D}_{z\circ r}^{-1} + O(1/n^2).
 \end{aligned}$$

Therefore as $n \rightarrow \infty$ the off-diagonal blocks of \mathbf{C}^{-1} will be $O(1/n^2)$ while the (τ, τ) diagonal block is

$$\frac{1}{z_\tau}\mathbf{I}_{r_\tau} - \frac{1}{z_\tau r_\tau}\mathbf{1}_{r_\tau \times r_\tau} + O(1/n^2)$$

as desired.

Using Lemma B.4, we can analyse the column sums of⁴⁶ $\mathbf{C}_i(0)^{-1}\mathbf{M}\mathbf{C}_i(0)^{-1}$. In more detail, we use the formula of the lemma to estimate both copies of $\mathbf{C}_i(0)^{-1}$, and then expand this to write an expression for any column sum of $\mathbf{C}_i(0)^{-1}\mathbf{M}\mathbf{C}_i(0)^{-1}$. It follows straightforwardly from this calculation that all these column sums are $O(1/n)$ whenever all entries of \mathbf{M} are in $[-1, 1]$.

We can bound the higher-order terms in the Taylor expansion by the same technique: by differentiating equation (B.1) repeatedly in x , we obtain an expression for the k th derivative in terms of $\mathbf{C}_i(0)^{-1}$ and \mathbf{M} :

$$f^{(k)}(0) = k!\mathbf{C}_i(0)^{-1}\mathbf{M}\mathbf{C}_i(0)^{-1}\mathbf{M}\mathbf{C}_i(0)^{-1} \cdots \mathbf{M}\mathbf{C}_i(0)^{-1},$$

where \mathbf{M} appears k times in the product. By the same argument as above, we can show that the column sums of $f^{(k)}(0)/k!$ are bounded by a constant independent of n . The Taylor expansion is

$$f(\mathbf{A}) = \sum_k \frac{f^{(k)}(0)}{k!}x^k.$$

Since we take $\mathbf{A} \in \mathcal{W}_{\psi, \zeta}$, we can assume that x is $O(1/n)$. Because the column sums of each summand are bounded by a constant times x^k , the column sums of $f(\mathbf{A})$ are bounded by a constant.

Finally, because the variation in the column sums is $O(1/n)$ and the weights are proportional to the column sums, each weight varies by at most a multiplicative factor of γ_1/n for some γ_1 . We find that the first part of the lemma, which bounded the ratios between weights $W_{ij,t+1}/W_{i'j',t+1}$, holds.

Step 3. We complete the proof of Lemma B.1, which states that the covariance matrix of $r_{i,t+1}$ is in $\mathcal{W}_{\psi, \zeta'}$. Recall that $\zeta' = \lambda/n$ for some constant n , so we are showing that if the covariance matrix of the $r_{i,t}$ is in a neighbourhood $\mathcal{W}_{\psi, \zeta}$, then the covariance matrix in the next period is in a somewhat larger neighbourhood $\mathcal{W}_{\psi, \zeta'}$. The remainder of the argument then follows by the same arguments as in the proof of the first part of the lemma: we now bound the change in time- $(t+2)$ weights as we vary the covariances of time- $(t+1)$ social signals within this neighbourhood.

46. Recall we wrote $A(x) = A(0) + x\mathbf{M}$, and in (B.1) we expressed the derivative of f in x in terms of the matrix we exhibit here.

Recall that we decomposed each covariance $\text{Cov}(r_{i,t} - \theta_{t-1}, r_{j,t} - \theta_{t-1}) = \psi_{kk'} + \zeta_{ij}$ into a term $\psi_{kk'}$ depending only on the types of the two agents and a term ζ_{ij} , and similarly for variances. To show the covariance matrix is contained in $\mathcal{W}_{\psi, \zeta'}$, we bound each of these terms suitably.

We begin with ζ_{ij} (and ζ_i). We can write

$$r_{i,t+1} = \sum_j \frac{W_{ij,t+1}}{1 - w_{i,t+1}^s} \rho a_{i,t} = \sum_j \frac{W_{ij,t+1}}{1 - w_{i,t+1}^s} \rho (w_{j,t}^s s_{j,t} + (1 - w_{j,t}^s) r_{j,t}).$$

By the first part of the lemma, the ratio between any two weights (both of the form $W_{ij,t+1}$, $w_{i,t+1}^s$, or $w_{j,t}^s$) corresponding to pairs of agents of the same types is in $[1 - \gamma_1/n, 1 + \gamma_1/n]$ for a constant γ_1 . We can use this to bound the variation in covariances of $r_{i,t+1}$ within types by ζ' : we take the covariance of $r_{i,t+1}$ and $r_{j,t+1}$ using the expansion above and then bound the resulting summation by bounding all coefficients.

Next we bound $\psi_{kk'}$ (and ψ_k). It is sufficient to show that $\text{Var}(r_{i,t+1} - \theta_t)$ is at most ψ . To do so, we will give an estimator of θ_t with variance less than β/n , and this will imply $\text{Var}(r_{i,t+1} - \theta_t) < \beta/n = \psi$ (recall $r_{i,t+1}$ is the estimate of θ_t given agent i 's social observations in period $t + 1$). Since this bounds all the variance terms by ψ , the covariance terms will also be bounded by ψ in absolute value.

Fix an agent i of network type k and consider some network type k' such that $p_{kk'} > 0$. Then there exists two signal types, which we call A and B , such that i observes $\Omega(n)$ agents of each of these signal types in G_n^k .⁴⁷ The basic idea will be that we can approximate θ_t well by taking a linear combination of the average of observed agents of network type k and signal type A and the average of observed agents of network type k and signal type B .

In more detail, let $N_{i,A}$ be the set of agents of type A in network type k observed by i and $N_{i,B}$ be the set of agents of type B in network type k observed by i . Then fixing some agent j_0 of network type k ,

$$\frac{1}{|N_{i,A}|} \sum_{j \in N_{i,A}} a_{j,t-1} = \frac{\sigma_A^{-2}}{1 + \sigma_A^{-2}} \theta_t + \frac{1}{1 + \sigma_A^{-2}} r_{j_0,t-1} + \text{noise}$$

where the noise term has variance of order $1/n$ and depends on signal noise, variation in $r_{j,t}$, and variation in weights. These bounds on the noise term follow from the assumption that the covariance matrix of the $r_{i,t}$ is in a neighbourhood $\mathcal{W}_{\psi, \zeta}$ and our analysis of variation in weights. Similarly

$$\frac{1}{|N_{i,B}|} \sum_{j \in N_{i,B}} a_{j,t-1} = \frac{\sigma_B^{-2}}{1 + \sigma_B^{-2}} \theta_t + \frac{1}{1 + \sigma_B^{-2}} r_{j_0,t-1} + \text{noise}$$

where the noise term has the same properties. Because $\sigma_A^2 \neq \sigma_B^2$, we can write θ_t as a linear combination of these two averages with coefficients independent of n up to a noise term of order $1/n$. We can choose β large enough such that this noise term has variance most β/n for all n sufficiently large. This completes the proof of Lemma B.1.

Step 4. We now give the two-step version of Lemma B.3.

47. We use the notation $\Omega(n)$ to mean greater than Cn for some constant $C > 0$ when n is large.

Lemma B.5. *Suppose that in period t the matrix $\mathbf{A} = \mathbf{A}_t$ of covariances of social signals satisfies $\mathbf{A} \in \mathcal{W}_{\psi, \zeta}$ and all agents are optimising in periods $t+1$ and $t+2$. Then there is a γ such that for all n sufficiently large,*

$$\frac{W_{ij,t+2}}{W_{i'j',t+2}} \in \left[1 - \frac{\gamma}{n}, 1 + \frac{\gamma}{n} \right].$$

whenever i and i' have the same network and signal types and j and j' have the same network and signal types.

Given what we established about covariances in Step 3, the lemma follows by the same argument as the proof of Lemma B.3.

Step 5. Now that Lemma B.5 is proved, we can apply it to show that $\tilde{\Phi}^2(\mathcal{W}_{\psi, \zeta}) \subset \mathcal{W}_{\psi, \zeta}$.

We will do this by first writing the time- $(t+2)$ behaviour in terms of agents' time- t observations (Step 5(a)), which comes from applying $\tilde{\Phi}$ twice. This gives a formula that can be used for bounding the covariances⁴⁸ of time- $(t+2)$ actions in terms of covariances of time- t actions. Step 5(b) then applies this formula to show we can take ζ_{ij} and ζ_i to be sufficiently small. (Recall the notation introduced in Section B.1 above.) We split our expression for $r_{i,t+2}$ into several groups of terms and show that the contribution of each group of terms depends only on agents' types up to a small noise term. Step 5(c) notes that we can also take $\psi_{kk'}$ and ψ_k to be sufficiently small.

Step 5(a): We calculate:

$$\begin{aligned} r_{i,t+2} &= \sum_j \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} \rho a_{j,t+1} \\ &= \rho \left(\sum_j \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} w_{j,t+1}^s s_{j,t+1} + \sum_{j,j'} \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} W_{jj',t+1} \rho a_{j',t} \right) \\ &= \rho \left(\sum_j \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} w_{j,t+1}^s s_{j,t+1} + \rho \left(\sum_{j,j'} \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} W_{jj',t+1} w_{j',t}^s s_{j',t} \right. \right. \\ &\quad \left. \left. + \sum_{j,j'} \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} W_{jj',t+1} (1 - w_{j',t}^s) r_{j',t} \right) \right). \end{aligned}$$

Let $h_{ij',t}$ be the coefficient on $r_{j',t}$ in this expansion of $r_{i,t+2}$. Explicitly,

$$h_{ij',t} = \sum_j \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} W_{jj',t+1} (1 - w_{j',t}^s). \quad (\text{B.3})$$

The coefficient $h_{ij',t}$ adds up the influence of $r_{j',t}$ on $r_{i,t+2}$ over all paths of length two.

First, we establish a lemma about how much these weights vary.

Lemma B.6. *There exists γ such that for n sufficiently large, when i and i' have the same network types and j' and j'' have the same network and signal types, the ratio $h_{ij',t} / h_{ij'',t}$ is in $[1 - \gamma/n, 1 + \gamma/n]$.*

48. We take this term to refer to variances, as well.

Proof. Fix i and j' . For each network type k'' and signal type τ , consider the number of agents j of network type k'' and signal type τ who are observed by i and who observe j' . This number varies by at most a factor ζ^2 as we vary i and j' , preserving signal and network types. For each such j , the contribution to $h_{ij',t}$ due to weight on that agent's action is (recalling (B.3))

$$\frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} W_{jj',t+1} (1 - w_{j',t}^s).$$

By applying Lemma B.3 repeatedly, we can choose γ_1 such that each of these contributions varies by at most a factor of γ_1/n as we change i in G_k and j' in $G_{k'}$. Thus, $h_{ij',t}$ is a sum of terms which vary by at most a multiplicative factor of γ_1/n as we vary i and j' preserving signal and network types. If we can show that the sum of the absolute values of these terms is bounded, then it will follow that $h_{ij',t}$ varies by at most a multiplicative factor of γ/n for some n . This bound on the sum of absolute values follows from the calculation of weights in the proof of Lemma B.3.

Step 5(b): We first show that fixing the values of $\psi_{kk'}$ and ψ_k in period t , the variation in the covariances $\text{Cov}(r_{i,t+2} - \theta_{t+1}, r_{i',t+2} - \theta_{t+1})$ of these terms as we vary i and i' over network types is not larger than ζ . From the formula above, we observe that we can decompose $r_{i,t+2} - \theta_{t+1}$ as a linear combination of three mutually independent groups of terms:

- (i) signal error terms $\eta_{j,t+1}$ and $\eta_{j',t}$;
- (ii) the errors $r_{j',t} - \theta_t$ in the social signals from period t ; and
- (iii) changes in state v_t and v_{t+1} between periods t and $t + 2$.

Note that the terms $r_{j',t} - \theta_t$ are linear combinations of older signal errors and changes in the state. We bound each of the three groups in turn:

(i) Signal errors: We first consider the contribution of signal errors. When i and i' are distinct, the number of such terms is close to its expected value because we are conditioning on the events E_2 and E_4 defined in Section B.1. Moreover, the weights are close to their expected values by Step 2, so the variation is bounded suitably. When i and i' are equal, we use the facts that the weights are close to their expected values and the variance of an average of $\Omega(n)$ signals is small.

(ii) Social signals: We now consider terms $r_{j',t} - \theta_t$, which correspond to the third summand in our expression for $r_{i,t+2}$. Since we will analyse the weight on v_t below, it is sufficient to study the terms $r_{j',t} - \theta_{t-1}$.

By Lemma B.6, the coefficients placed on $r_{j',t}$ by i and on $r_{j'',t}$ by i' vary by a factor of at most $2\gamma/n$. Moreover, the absolute value of each of these covariances is bounded above by ψ and the variation in these terms is bounded above by ζ . We conclude that the variation from these terms has order $1/n^2$.

(iii) Innovations: Finally, we consider the contribution of the innovations v_t and v_{t+1} . We treat v_{t+1} first. We must show that any two agents of the same types place the same weight on the innovation v_{t+1} (up to an error of order $1/n^2$). This will imply that the contributions of timing to the covariances $\text{Cov}(r_{i,t+2} - \theta_{t+1}, r_{i',t+2} - \theta_{t+1})$ can be expressed as a term that can be included in the relevant $\psi_{kk'}$ and a lower-order term which can be included in $\zeta_{ii'}$.

The weight an agent places on v_{t+1} is equal to the weight she places on signals from period $t + 1$. So this is equivalent to showing that the total weight

$$\rho \sum_j \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} w_{j,t+1}^s$$

agent i places on period $t + 1$ depends only on the network type k of agent i and $O_p(1/n^2)$ terms. We will first show the average weight placed on time- $(t + 1)$ signals by agents of each signal type depends only on k . We will then show that the total weights on agents of each signal type do not depend on n .

Suppose for simplicity here that there are two signal types A and B ; the general case is the same. We can split the sum from the previous paragraph into the subgroups of agents with signal types A and B :

$$\rho \sum_{j:\sigma_j^2=\sigma_A^2} \frac{W_{ij,t+2}}{1-w_{i,t+2}^s} w_{j,t+1}^s + \rho \sum_{j:\sigma_j^2=\sigma_B^2} \frac{W_{ij,t+2}}{1-w_{i,t+2}^s} w_{j,t+1}^s.$$

Letting $W_i^A = \sum_{j:\sigma_j^2=\sigma_A^2} W_{ij,t+2}/(1-w_{i,t+2}^s)$ be the total weight placed on agents with signal type A and similarly for signal type B , we can rewrite this as:

$$W_i^A \rho \sum_{j:\sigma_j^2=\sigma_A^2} \frac{W_{ij,t+2}}{W_i^A(1-w_{i,t+2}^s)} w_{j,t+1}^s + W_i^B \rho \sum_{j:\sigma_j^2=\sigma_B^2} \frac{W_{ij,t+2}}{W_i^B(1-w_{i,t+2}^s)} w_{j,t+1}^s.$$

The coefficients $W_{ij,t+2}/W_i^A(1-w_{i,t+2}^s)$ in the first sum now sum to one, and similarly for the second. We want to check that the first sum $\sum_{j:\sigma_j^2=\sigma_A^2} (W_{ij,t+2}/W_i^A(1-w_{i,t+2}^s)) w_{j,t+1}^s$ does not depend on k , and the second sum is similar.

For each j in group A ,

$$w_{j,t+1}^s = \frac{\sigma_A^{-2}}{\sigma_A^{-2} + (\rho^2 \kappa_{j,t+1} + 1)^{-1}},$$

where we define $\kappa_{j,t+1}^2 = \text{Var}(r_{j,t+1} - \theta_t)$ to be the error variance of the social signal. Because $\kappa_{j,t+1}$ is close to zero, we can approximate $w_{j,t+1}^s$ locally as a linear function $\mu_1 \kappa_{j,t+1} + \mu_2$ where $\mu_1 < 1$ (up to order $1/n^2$ terms).

So we can write the sum of interest as

$$\sum_{j:\sigma_j^2=\sigma_A^2} \frac{W_{ij,t+2}}{W_i^A(1-w_{i,t+2}^s)} \left(\mu_1 \sum_{j',j''} W_{jj',t+1} W_{jj'',t+1} (\rho^2 \mathbf{V}_{j'j'',t} + 1) + \mu_2 \right).$$

By Lemma B.3, the weights vary by at most a multiplicative factor contained in $[1 - \gamma/n, 1 + \gamma/n]$. The number of paths from i to j' passing through agents of any network type k'' and any signal type is close to its expected value (which depends only on i 's network type), and the weight on each path depends only on the types involved up to a factor in $[1 - \gamma/n, 1 + \gamma/n]$. The variation in $\mathbf{V}_{j'j'',t}$ consists of terms of the form $\psi_{kk''}$, $\psi_{k'}$, and $\zeta_{j'j''}$, all of which are $O_p(1/n)$, and terms from signal errors $\eta_{j',t}$. The signal errors only contribute when $j = j'$, and so only contribute to a fraction of the summands of order $1/n$. So we can conclude the total variation in this sum as we change i within the network type k has order $1/n^2$.

Now that we know each the average weight on private signals of the observed agents of each signal type depends only on k , it remains to check that W_i^A and W_i^B only depend on k . The coefficients W_i^A and W_i^B are the optimal weights on the group averages

$$\sum_{j:\sigma_j^2=\sigma_A^2} \frac{W_{ij,t+2}}{W_i^A(1-w_{i,t+2}^s)} \rho a_{j,t+1} \quad \text{and} \quad \sum_{j:\sigma_j^2=\sigma_B^2} \frac{W_{ij,t+2}}{W_i^B(1-w_{i,t+2}^s)} \rho a_{j,t+1},$$

so we need to show that the variances and covariance of these two terms depend only on k . We check the variance of the first sum: we can expand

$$\sum_{\sigma_j^2 = \sigma_A^2} \frac{W_{ij,t+2}}{W_i^A(1 - w_{i,t+2}^s)} \rho a_{j,t+1} = \sum_{\sigma_j^2 = \sigma_A^2} \frac{W_{ij,t+2}}{W_i^A(1 - w_{i,t+2}^s)} \rho (w_{j,t+1}^s s_{j,t+1} + (1 - w_{j,t+1}^s) r_{j,t+1}).$$

We can again bound the signal errors and social signals as in the previous parts of this proof, and show that the variance of this term depends only on k up to error terms that are $O_p(1/n^2)$. The second variance and covariance are similar, so W_i^A and W_i^B depend only on k up to error terms that are $O_p(1/n^2)$.

This takes care of the innovation v_{t+1} . Because we have included any innovations prior to v_t in the social signals $r_{j',t}$, to complete Step 5(b) we need only show the weight on v_t depends only on the network type k of an agent.

The analysis is a simpler version of the analysis of the weight on v_{t+1} . It is sufficient to show the total weight placed on period t social signals depends only on the network type of k of an agent i . This weight is equal to

$$\rho^2 \sum_{j,j'} \frac{W_{ij,t+2}}{1 - w_{i,t+2}^s} \cdot W_{jj',t+1} \cdot (1 - w_{j',t}^s).$$

As in the v_{t+1} case, we can approximate $(1 - w_{j',t}^s)$ as a linear function of $\kappa_{j',t}$ up to $O_p(1/n^2)$ terms. Because the number of paths to each agent j' through a given type and the weights on each such path cannot vary too much within types, the same argument shows that this sum depends only on k , up to error terms that are $O_p(1/n^2)$. Thus Step 5(b) is complete.

Step 5(c): The final step is to verify that we can take $\psi_{kk'}$ and ψ_k to be smaller than ψ . It is sufficient to show that the variance $\text{Var}(r_{i,t+2} - \theta_{t+1})$ of each social signal about θ_{t+1} is at most ψ . The proof is the same as in Step 2(b).

Acknowledgments. We gratefully acknowledge funding from Pershing Square Fund for Research on the Foundations of Human Behaviour at Harvard, the Rockefeller Foundation (Dasaratha), and the National Science Foundation under grants SES-1847860 and SES-1629446 (Golub), as well as the hospitality of the economics department at the University of Pennsylvania during a key phase of this work. Hershdeep Chopra, Yu-Chi Hsieh, Yixi Jiang, Joey Feffer, and Rithvik Rao provided excellent research assistance. For valuable comments we are grateful to (in random order) Erik Madsen, Alireza Tahbaz-Salehi, Drew Fudenberg, Matt Elliott, Nageeb Ali, Tomasz Strzalecki, Alex Frankel, Yair Livne, Jeroen Swinkels, Margaret Meyer, Philipp Strack, Erik Eyster, Michael Powell, Michael Ostrovsky, Leeat Yariv, Andrea Galeotti, Eric Maskin, Elliot Lipnowski, Jeff Ely, Eddie Dekel, Annie Liang, Iosif Pinelis, Ozan Candogan, Ariel Rubinstein, Paul Goldsmith-Pinkham, Bob Wilson, Omer Tamuz, Jeff Zwiebel, Matthew O. Jackson, Matthew Rabin, and Andrzej Skrzypacz. We also thank Evan Sadler for important conversations early in the project, David Hirshleifer for detailed comments on a draft, and Rohan Dutta and Kevin He for helpful discussions. The paper was greatly improved by advice from four anonymous referees and the editor, Adam Szeidl. All errors are our own.

Supplementary Data

Supplementary data are available at the *Review of Economic Studies* online. The data and code underlying this article are available on Zenodo under: <https://dx.doi.org/10.5281/zenodo.6954517>.

REFERENCES

- ABBE, E. (2017), "Community detection and stochastic block models: recent developments", *The Journal of Machine Learning Research*, **18**, 6446–6531.
- ACEMOGLU, D., DAHLEH, M., LOBEL, I. and OZDAGLAR, A. (2011), "Bayesian learning in social networks", *The Review of Economic Studies*, **78**, 1201–1236.
- ACEMOGLU, D. and OZDAGLAR, A. (2011), "Opinion dynamics and learning in social networks", *Dynamic Games and Applications*, **1**, 3–49.

- ALATAS, V., BANERJEE, A., CHANDRASEKHAR, A. G., HANNA, R. and OLKEN, B. A. (2016), "Network structure and the aggregation of information: theory and evidence from Indonesia", *The American Economic Review*, **106**, 1663–1704.
- ANGELETOS, G.-M. and LA'O, J. (2010), "Noisy business cycles", *NBER Macroeconomics Annual*, **24**, 319–378.
- ANGELETOS, G.-M. and LIAN, C. (2018), "Forward guidance without common knowledge", *American Economic Review*, **108**, 2477–2512.
- ARIELI, I. and MUELLER-FRANK, M. (2019), "Multidimensional social learning", *The Review of Economic Studies*, **86**, 913–940.
- BABUS, A. and KONDOR, P. (2018), "Trading and information diffusion in over-the-counter markets", *Econometrica*, **86**, 1727–1769.
- BALA, V. and GOYAL, S. (1998), "Learning from neighbours", *The Review of Economic Studies*, **65**, 595–621.
- BANERJEE, A., CHANDRASEKHAR, A. G., DUFLO, E. and JACKSON, M. O. (2013), *The Diffusion of Microfinance*, Harvard Database. <https://doi.org/10.7910/DVN/U3BIHX>.
- BANERJEE, A. and FUDENBERG, D. (2004), "Word-of-mouth learning", *Games and economic behavior*, **46**, 1–22.
- DASARATHA, K., GOLUB, B. and HAK, N. (2018), "Bayesian Social Learning in a Dynamic Environment", arXiv:1801.02042v1.
- DASARATHA, K. and HE, K. (2021), "An experiment on network density and sequential learning", *Games and Economic Behavior*, **128**, 182–192.
- DEGROOT, M. H. (1974), "Reaching a consensus", *Journal of the American Statistical Association*, **69**, 118–121.
- DEMARZO, P., VAYANOS, D. and ZWEIBEL, J. (2003), "Persuasion bias, social influence, and unidimensional opinions", *The Quarterly Journal of Economics*, **118**, 909–968.
- EATON, M. L. (1983), *Multivariate Statistics: A Vector Space Approach* (New York, NY: John Wiley & Sons, Inc.).
- ENKE, B. and ZIMMERMANN, F. (2019), "Correlation neglect in belief formation", *The Review of Economic Studies*, **86**, 313–332.
- EYSTER, E. and RABIN, M. (2010), "Naive herding in rich-information settings", *American Economic Journal: Microeconomics*, **2**, 221–243.
- EYSTER, E. and RABIN, M. (2014), "Extensive imitation is irrational and harmful", *Quarterly Journal of Economics*, **129**, 1861–1898.
- EYSTER, E., RABIN, M. and WEIZSACKER, G. (2015), "An experiment on social mislearning", available at SSRN 2704746.
- FRIEDKIN, N. E. and JOHNSEN, E. C. (1997), "Social positions in influence networks", *Social Networks*, **19**, 209–222.
- FRONGILLO, R. M., SCHOENEBECK, G. and TAMUZ, O. (2011), "Social learning in a changing world", in *International Workshop on Internet and Network Economics* (Springer) 146–157.
- GOEREE, J. K., PALFREY, T. R. and ROGERS, B. W. (2006), "Social learning with private and common values", *Economic Theory*, **28**, 245–264. AQ5
- GOLUB, B. and JACKSON, M. (2010), "Naïve learning in social networks and the wisdom of crowds", *American Economic Journal: Microeconomics*, **2**, 112–149.
- GOLUB, B. and SADLER, E. (2016), "Learning in social networks", in Bramoullé, Y., Galeotti, A., Rogers, B. and Rogers, B. (eds) *The Oxford Handbook of the Economics of Networks*, chap. 19 (Oxford University Press) 504–542. AQ6
- HAREL, M., MOSSEL, E., STRACK, P. and TAMUZ, O. (2021), "Rational groupthink", *The Quarterly Journal of Economics*, **136**, 621–668.
- HAYEK, F. A. (1945), "The use of knowledge in society", *The American Economic Review*, **35**, 519–530.
- HOLLAND, P. W., LASKEY, K. B. and LEINHARDT, S. (1983), "Stochastic blockmodels: first steps", *Social Networks*, **5**, 109–137.
- JACKSON, M. O. (2010), *Social and Economic Networks* (Princeton University Press). AQ7
- JADBABAIE, A., MOLAVI, P., SANDRONI, A. and TAHBAZ-SALEHI, A. (2012), "Non-Bayesian social learning", *Games and Economic Behavior*, **76**, 210–225.
- JANSON, S. (2011), "Probability asymptotics: notes on notation", arXiv:1108.3924.
- LAMBERT, N. S., OSTROVSKY, M. and PANOV, M. (2018), "Strategic trading in informationally complex environments", *Econometrica*, **86**, 1119–1157.
- LOBEL, I. and SADLER, E. (2015), "Information diffusion in networks through social learning", *Theoretical Economics*, **10**, 807–851.
- LOBEL, I. and SADLER, E. (2016), "Preferences, homophily, and social learning", *Operations Research*, **64**, 564–584.
- MOLAVI, P. (2019), "Macroeconomics with learning and misspecification: a general theory and applications" (Working Paper, Kellogg School of Management, Evanston, IL).
- MOLAVI, P., TAHBAZ-SALEHI, A. and JADBABAIE, A. (2018), "A theory of non-Bayesian social learning", *Econometrica*, **86**, 445–490.
- MOSCARINI, G., OTTAVIANI, M. and SMITH, L. (1998), "Social learning in a changing world", *Economic Theory*, **11**, 657–665.
- MOSSEL, E., MUELLER-FRANK, M., SLY, A. and TAMUZ, O. (2020), "Social learning equilibria", *Econometrica*, **88**, 1235–1267.
- MOSSEL, E. and TAMUZ, O. (2017), "Opinion exchange dynamics", *Probability Surveys*, **14**, 155–204.
- MYATT, D. and WALLACE, C. (2017), "Information acquisition and use by networked players" (University of Warwick, Department of Economics). CRETA Discussion Paper Series (32), Available at <http://wrap.warwick.ac.uk/90449/>.

- PINELIS, I. (2018), "Inverse of matrix with blocks of ones", *MathOverflow*. <https://mathoverflow.net/q/296933> (version: 2018-04-04).
- SETHI, R. and YILDIZ, M. (2012), "Public disagreement", *American Economic Journal: Microeconomics*, **4**, 57–95.
- SETHI, R. and YILDIZ, M. (2016), "Communication with unknown perspectives", *Econometrica*, **84**, 2029–2069.
- SETHI, R. and YILDIZ, M. (2019), "Culture and communication", available at SSRN 3263560.
- SGROI, D. (2002), "Optimizing information in the herd: guinea pigs, profits, and welfare", *Games and Economic Behavior*, **39**, 137–166.
- SHAHRAMPOUR, S., RAKHLIN, S. and JADBABAIE, A. (2013), "Online learning of dynamic parameters in social networks", *Advances in Neural Information Processing Systems*.
- VIVES, X. (1993), "How fast do rational agents learn?", *The Review of Economic Studies*, **60**, 329–347. AQ8
- WEIZSÄCKER, G. (2010), "Do we follow others when we should? A simple test of rational expectations", *The American Economic Review*, **100**, 2340–2360.
- WOLITZKY, A. (2018), "Learning from others' outcomes", *American Economic Review*, **108**, 2763–2801.